

# Calibrated Submanifolds

Jason D. Lotay  
University College London

## Abstract

We provide an introduction to the theory of calibrated submanifolds through the key examples related with special holonomy. We focus on calibrated geometry in Calabi–Yau,  $G_2$  and  $\text{Spin}(7)$  manifolds, and describe fundamental results and techniques in the field.

## Contents

1	Minimal submanifolds	2
2	Introduction to calibrations	4
3	Complex submanifolds	5
4	Special Lagrangians	6
5	Constructing calibrated submanifolds	9
6	Associative and coassociative submanifolds	13
7	Cayley submanifolds	17
8	The angle theorem	19

## Introduction

A key aspect of mathematics is the study of variational problems. These can vary from the purely analytic to the very geometric. A classic geometric example is the study of geodesics, which are critical points for the length functional on curves. As we know, understanding the geodesics of a given Riemannian manifold allows us to understand some of the ambient geometry, for example the curvature. The higher dimensional analogue would be to study critical points for the volume functional, and we would hope (and it indeed turns out to be the case) that these critical points, called *minimal submanifolds*, encode crucial aspects of the geometry of the manifold.

Just like the geodesic equation, we would expect (and it is true) that minimal submanifolds are defined by a (nonlinear) second order partial differential equation. Such equations are very difficult to solve in general, so a key idea is to find a special class of minimal submanifolds, called *calibrated submanifolds*, which are instead defined by a first order partial differential equation. The definition of calibrated submanifolds is motivated by the properties of complex submanifolds in Kähler manifolds, and turns out to be useful in finding minimizers for the volume functional rather than just critical points. However, finding examples outside the classical complex setting turns out to be difficult, leading to important methods coming from a variety of sources, as well as motivating the study of the deformation theory of these objects.

Calibrated submanifolds naturally arise when the ambient manifold has *special holonomy*, including holonomy  $G_2$ . In this situation, we would hope that the calibrated submanifolds encode even more, finer, information about the ambient manifold, potentially leading to the construction of new invariants. In this setting, there is also a relationship between calibrated submanifolds and gauge theory: specifically,

connections whose curvature satisfies a natural constraint determined by the special holonomy group (so-called *instantons*). For these reasons, calibrated submanifolds form a hot topic in current research, especially in the  $G_2$  setting.

**Note** These notes are primarily based on a lecture course the author gave at the LMS–CMI Research School “An Invitation to Geometry and Topology via  $G_2$ ” at Imperial College London in July 2014.

## 1 Minimal submanifolds

We start by analysing the submanifolds which are critical points for the volume functional. Let  $N$  be a submanifold (without boundary) of a Riemannian manifold  $(M, g)$  and let  $F : N \times (-\epsilon, \epsilon) \rightarrow M$  be a variation of  $N$  with compact support; i.e.  $F = \text{Id}$  outside a compact subset  $\bar{S}$  of  $N$  with  $S$  open and  $F(p, 0) = p$  for all  $p \in N$ . The vector field  $X = \frac{\partial F}{\partial t}|_N$  is called the variation vector field (which will be zero outside of  $\bar{S}$ ). We then have the following definition.

**Definition 1.1.**  $N$  is *minimal* if  $\frac{d}{dt} \text{Vol}(F(S, t))|_{t=0} = 0$  for all variations  $F$  with compact support  $\bar{S}$  (depending on  $F$ ).

**Remark.** Notice that we do not ask for  $N$  to minimize volume: it is only stationary for the volume. It could even be a maximum!

**Example.** A plane in  $\mathbb{R}^n$  is minimal since any small variation will have larger volume.

**Example.** Geodesics are locally length minimizing, so geodesics are minimal. However, as an example, the equator in  $S^2$  is minimal but not length minimizing since we can deform it to a shorter line of latitude.

For simplicity let us suppose that  $N$  is compact. We wish to calculate  $\frac{d}{dt} \text{Vol}(F(N, t))|_{t=0}$ . Given local coordinates  $x_i$  on  $N$  we know that

$$\text{Vol}(F(N, t)) = \int_N \sqrt{\det \left( g \left( \frac{\partial F}{\partial x_i}, \frac{\partial F}{\partial x_j} \right) \right)} \text{vol}_N.$$

Let  $p \in N$  and choose our coordinates  $x_i$  to be normal coordinates at  $p$ : i.e. so that  $\frac{\partial F}{\partial x_i}(p, t) = e_i(t)$  satisfy  $g(e_i(0), e_j(0)) = \delta_{ij}$ . If  $g_{ij}(t) = g(e_i(t), e_j(t))$  and  $(g^{ij}(t))$  denotes the inverse of the matrix  $(g_{ij}(t))$  then we know that

$$\frac{d}{dt} \sqrt{\det(g_{ij}(t))}|_{t=0} = \frac{1}{2} \frac{\sum_{i,j} g^{ij}(t) g'_{ij}(t)}{\sqrt{\det(g_{ij}(t))}}|_{t=0} = \frac{1}{2} \sum_i g'_{ii}(0).$$

Now, if we let  $\nabla$  denote the Levi-Civita connection of  $g$ , then

$$\begin{aligned} \frac{1}{2} \sum_i g'_{ii}(0) &= \frac{1}{2} \sum_i \frac{d}{dt} g \left( \frac{\partial F}{\partial x_i}, \frac{\partial F}{\partial x_i} \right) |_{t=0} \\ &= \sum_i g(\nabla_X e_i, e_i) \\ &= \sum_i g(\nabla_{e_i} X, e_i) = \text{div}_N(X) \end{aligned}$$

since  $[X, e_i] = 0$  (i.e. the  $t$  and  $x_i$  derivatives commute). Moreover, we see that

$$\text{div}_N(X) = \sum_i g(\nabla_{e_i} X, e_i) = \text{div}_N(X^T) - \sum_i g(X^\perp, \nabla_{e_i} e_i) = \text{div}_N(X^T) - g(X, H)$$

(since  $\nabla_{e_i}(g(X^\perp, e_i)) = 0$ ) where  $^T$  and  $^\perp$  denote the tangential and normal parts and

$$H = \sum_i \nabla_{e_i}^\perp e_i$$

is the *mean curvature vector*. Overall we have the following.

**Theorem 1.2.** *The first variation formula is*

$$\frac{d}{dt} \text{Vol}(F(N, t))|_{t=0} = \int_N \text{div}_N(X) \text{vol}_N = - \int_N g(X, H) \text{vol}_N.$$

**Remark.** The  $\text{div}_N(X^T)$  term does not appear in the first variation formula because its integral vanishes by the divergence theorem as  $N$  is compact without boundary. In general, it will still vanish since we assume for our variations that there exists a compact submanifold of  $N$  with boundary which contains the support of  $X^T$  and so that  $X^T$  vanishes on the boundary.

We deduce the following.

**Definition 1.3.**  $N$  is a *minimal submanifold* if and only if  $H = 0$ .

The equation  $H = 0$  is a *second order nonlinear PDE*. We can see this explicitly in the following simple case. For a function  $f : U \subseteq \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  where  $\bar{U}$  is compact, we see that if  $N = \text{Graph}(f) \subseteq \mathbb{R}^n$  then the volume of  $N$  is given by

$$\text{Vol}(N) = \int_U \sqrt{1 + |\nabla f|^2} \text{vol}_U.$$

Any sufficiently small variation can be written  $F(N, t) = \text{Graph}(f + th)$  for some  $h : U \rightarrow \mathbb{R}$ , so we can compute

$$\begin{aligned} \frac{d}{dt} \text{Vol}(F(N, t))|_{t=0} &= \frac{d}{dt}|_{t=0} \int_U \sqrt{1 + |\nabla f + t\nabla h|^2} \text{vol}_U \\ &= \int_U \frac{d}{dt}|_{t=0} \sqrt{1 + |\nabla f|^2 + 2t\langle \nabla f, \nabla h \rangle + t^2|\nabla h|^2} \text{vol}_U \\ &= \int_U \frac{\langle \nabla f, \nabla h \rangle}{\sqrt{1 + |\nabla f|^2}} \text{vol}_U \\ &= - \int_U h \text{div} \left( \frac{\nabla f}{\sqrt{1 + |\nabla f|^2}} \right) \text{vol}_U. \end{aligned}$$

We therefore see that  $N$  is minimal if and only if this vanishes for all  $h$ . Hence,  $\text{Graph}(f)$  is minimal in  $\mathbb{R}^n$  if and only if

$$\text{div} \left( \frac{\nabla f}{\sqrt{1 + |\nabla f|^2}} \right) = 0.$$

We see that we can write this equation as  $\Delta f + Q(\nabla f, \nabla^2 f) = 0$  where  $Q$  consists of nonlinear terms (but linear in  $\nabla^2 f$ ). Hence, if we linearise this equation we just get  $\Delta f = 0$ , so  $f$  is harmonic. In other words, the minimal submanifold equation is a nonlinear equation whose linearisation is just Laplace's equation: this is an example of a nonlinear *elliptic* PDE, which we shall discuss further later.

**Example.** A plane in  $\mathbb{R}^n$  is trivially minimal because if  $X, Y$  are any vector fields on the plane then  $\nabla_X^\perp Y = 0$  as the second fundamental form of a plane is zero.

**Example.** For curves  $\gamma$ ,  $H = 0$  is equivalent to the geodesic equation  $\nabla_{\dot{\gamma}} \dot{\gamma} = 0$ .

The most studied minimal submanifolds (other than geodesics) are minimal surfaces in  $\mathbb{R}^3$ , since here the equation  $H = 0$  becomes a scalar equation on a surface, which is the simplest to analyse. In general we would have a system of equations, which is more difficult to study.

**Example.** The helicoid  $M = \{(t \cos s, t \sin s, s) \in \mathbb{R}^3 : s, t \in \mathbb{R}\}$  is a complete embedded minimal surface, discovered by Meusnier in 1776.

**Example.** The catenoid  $M = \{(\cos t \cos s, \cos t \sin s, t) \in \mathbb{R}^3 : s, t \in \mathbb{R}\}$  is a complete embedded minimal surface, discovered by Euler in 1744 and shown to be minimal by Meusnier in 1776. The catenoid is another explicit example which is a critical point for volume but not minimizing.

In fact the helicoid and the catenoid are locally isometric, and there is a 1-parameter family of locally isometric minimal surfaces deforming between the catenoid and helicoid: see, for example, [18, Theorem 16.5] for details.

It took about 70 years to find the next minimal surface, but now we know many examples of minimal surfaces in  $\mathbb{R}^3$ , as well as in other spaces by studying the nonlinear elliptic PDE given by the minimal surface equation. The amount of literature in the area is vast, with key results including the proofs of the Lawson [1], Willmore [63] and Yau [29, 64, 77] Conjectures, and minimal surfaces have applications to major problems in geometry including the Positive Mass Theorem [75, 76], Penrose Inequality [24] and Poincaré Conjecture [74].

## 2 Introduction to calibrations

As we have seen, minimal submanifolds are extremely important. However there are two key issues.

- Minimal submanifolds are defined by a second order nonlinear PDE system – therefore they are hard to analyse.
- Minimal submanifolds are only critical points for the volume functional, but we are often interested in minima for the volume functional – we need a way to determine when this occurs.

We can help resolve these issues using the notion of calibration and calibrated submanifolds, introduced by Harvey–Lawson [20] in 1982.

**Definition 2.1.** A differential  $k$ -form  $\eta$  on a Riemannian manifold  $(M, g)$  is a *calibration* if

- $d\eta = 0$  and
- $\eta(e_1, \dots, e_k) \leq 1$  for all unit tangent vectors  $e_1, \dots, e_k$  on  $M$ .

**Example.** Any non-zero form with constant coefficients on  $\mathbb{R}^n$  can be rescaled so that it is a calibration with at least one plane where equality holds.

This example shows that there are many calibrations  $\eta$ , but the interesting question is: for which oriented planes  $P = \text{Span}\{e_1, \dots, e_k\}$  does  $\eta(e_1, \dots, e_k) = 1$ ? More importantly, can we find submanifolds  $N$  so that this equality holds on each tangent space? This motivates the next definition.

**Definition 2.2.** Let  $\eta$  be a calibration  $k$ -form on  $(M, g)$ . An oriented  $k$ -dimensional submanifold  $N$  of  $(M, g)$  is *calibrated* by  $\eta$  if  $\eta|_N = \text{vol}_N$ , i.e. if for all  $p \in N$  we have  $\eta(e_1, \dots, e_k) = 1$  for an oriented orthonormal basis  $e_1, \dots, e_k$  for  $T_p N$ .

**Example.** Any oriented plane in  $\mathbb{R}^n$  is calibrated. If we change coordinates so that the plane  $P$  is  $\{x \in \mathbb{R}^n : x_{k+1} = \dots = x_n = 0\}$  (with the obvious orientation) then  $\eta = dx_1 \wedge \dots \wedge dx_k$  is a calibration and  $P$  is calibrated by  $\eta$ .

Notice that the calibrated condition is now an algebraic condition on the tangent vectors to  $N$ , so being calibrated is a *first order nonlinear PDE*. We shall motivate these definitions further later, but for now we make the following observation.

**Theorem 2.3.** *Let  $N$  be a calibrated submanifold. Then  $N$  is minimal and, moreover, if  $F$  is any variation with compact support  $\bar{S}$  then  $\text{Vol}(F(S, t)) \geq \text{Vol}(S)$ ; i.e.  $N$  is volume-minimizing. In particular, if  $N$  is compact then  $N$  is volume-minimizing in its homology class.*

*Proof.* Suppose that  $N$  is calibrated by  $\eta$  and suppose for simplicity that  $N$  is compact. We will show that  $N$  is homologically volume-minimizing.

Suppose that  $N'$  is homologous to  $N$ . Then there exists a compact manifold  $K$  with boundary  $-N \cup N'$  and, since  $d\eta = 0$ , we have by Stokes' Theorem that

$$0 = \int_K d\eta = \int_{N'} \eta - \int_N \eta.$$

We deduce that

$$\text{Vol}(N) = \int_N \eta = \int_{N'} \eta \leq \text{Vol}(N').$$

We then have the result by the definition of minimal submanifold.  $\square$

We conclude this introduction with the following elementary result.

**Proposition 2.4.** *There are no compact calibrated submanifolds in  $\mathbb{R}^n$ .*

*Proof.* Suppose that  $\eta$  is a calibration and  $N$  is compact and calibrated by  $\eta$ . Then  $d\eta = 0$  so by the Poincaré Lemma  $\eta = d\zeta$ , and hence

$$\text{Vol}(N) = \int_N \eta = \int_N d\zeta = 0$$

by Stokes' Theorem. □

Although there are many calibrations, having calibrated submanifolds greatly restricts the calibrations you want to consider. The calibrations which have calibrated submanifolds have special significance and there is a particular connection with special holonomy, due to the following observations.

Let  $G$  be the holonomy group of a Riemannian metric  $g$  on an  $n$ -manifold  $M$ . Then  $G$  acts on the  $k$ -forms on  $\mathbb{R}^n$ , so suppose that  $\eta_0$  is a  $G$ -invariant  $k$ -form. We can always rescale  $\eta_0$  so that  $\eta_0|_P \leq \text{vol}_P$  for all oriented  $k$ -planes  $P$  and equality holds for at least one  $P$ . Since  $\eta_0$  is  $G$ -invariant, if  $P$  is calibrated then so is  $\gamma \cdot P$  for any  $\gamma \in G$ , which usually means we have quite a few calibrated planes. We know by the *holonomy principle* (see, for example, [42, Proposition 2.5.2]) that we then get a parallel  $k$ -form  $\eta$  on  $M$  which is identified with  $\eta_0$  at every point. Since  $\nabla\eta = 0$ , we have  $d\eta = 0$  and hence  $\eta$  is a calibration. Moreover, we have a lot of calibrated tangent planes on  $M$ , so we can hope to find calibrated submanifolds.

### 3 Complex submanifolds

We would now like to address the question: where does the calibration condition come from? The answer is from *complex geometry*. On  $\mathbb{R}^{2n} = \mathbb{C}^n$  with coordinates  $z_j = x_j + iy_j$ , we have the complex structure  $J$  and the distinguished Kähler 2-form

$$\omega = \sum_{j=1}^n dx_j \wedge dy_j = \frac{i}{2} \sum_{j=1}^n dz_j \wedge d\bar{z}_j.$$

More generally we can work with a *Kähler manifold*  $(M, J, \omega)$ . Our first key result is the following.

**Theorem 3.1.** *On a Kähler manifold  $(M, J, \omega)$ ,  $\frac{\omega^k}{k!}$  is a calibration whose calibrated submanifolds are the complex  $k$ -dimensional submanifolds: i.e. submanifolds  $N$  such that  $J(T_p N) = T_p N$  for all  $p \in N$ .*

Since  $d\omega^k = k d\omega \wedge \omega^{k-1} = 0$ , Theorem 3.1 follows immediately from the following result.

**Theorem 3.2** (Wirtinger's inequality). *For any unit vectors  $e_1, \dots, e_{2k} \in \mathbb{C}^n$ ,*

$$\frac{\omega^k}{k!}(e_1, \dots, e_{2k}) \leq 1$$

*with equality if and only if  $\text{Span}\{e_1, \dots, e_{2k}\}$  is a complex  $k$ -plane in  $\mathbb{C}^n$ .*

Before proving this we make the following observation.

**Lemma 3.3.** *If  $\eta$  is a calibration and  $*\eta$  is closed then  $*\eta$  is a calibration. Moreover an oriented tangent plane  $P$  is calibrated by  $\eta$  if and only if there is an orientation on the orthogonal complement  $P^\perp$  so that it is calibrated by  $*\eta$ .*

*Proof.* Suppose that  $\eta$  is a calibration  $k$ -form on  $(M, g)$  with  $d*\eta = 0$ . Let  $p \in M$ . Take any  $n - k$  orthonormal tangent vectors  $e_{k+1}, \dots, e_n$  at  $p$ . Then there exist  $e_1, \dots, e_k \in T_p M$  so that  $\{e_1, \dots, e_n\}$  is an oriented orthonormal basis for  $T_p M$ . Since  $\{e_1, \dots, e_n\}$  is an oriented orthonormal basis, we can use the definition of the Hodge star to calculate

$$*\eta(e_{k+1}, \dots, e_n) = \eta(e_1, \dots, e_k) \leq 1.$$

Hence  $*\eta$  is a calibration by Definition 2.1. Moreover, the oriented plane  $P = \text{Span}\{e_{k+1}, \dots, e_n\}$  is calibrated by  $*\eta$  if and only if there is an orientation on  $\text{Span}\{e_1, \dots, e_k\} = P^\perp$  so that it is calibrated by  $\eta$ , since  $\eta(e_1, \dots, e_k) = \pm *\eta(e_{k+1}, \dots, e_n) = \pm 1$ . □

We can now prove Wirtinger's inequality.

*Proof of Theorem 3.2.* We see that  $|\frac{\omega^k}{k!}|^2 = \frac{n!}{k!(n-k)!}$  and  $\text{vol}_{\mathbb{C}^n} = \frac{\omega^n}{n!}$  so  $*\frac{\omega^k}{k!} = \frac{\omega^{n-k}}{(n-k)!}$ . Hence, by Lemma 3.3, it is enough to study the case where  $k \leq \frac{n}{2}$ .

Let  $P$  be any  $2k$ -plane in  $\mathbb{C}^n$  with  $2k \leq n$ . We shall find a canonical form for  $P$ . First consider  $\langle Ju, v \rangle$  for orthonormal vectors  $u, v \in P$ . This must have a maximum, so let  $\cos \theta_1 = \langle Ju, v \rangle$  be this maximum realised by some orthonormal vectors  $u, v \in P$ , where  $0 \leq \theta_1 \leq \frac{\pi}{2}$ .

Suppose that  $w \in P$  is a unit vector orthogonal to  $\text{Span}\{u, v\}$ , where  $\cos \theta_1 = \langle Ju, v \rangle$ . The function

$$f_w(\theta) = \langle Ju, \cos \theta v + \sin \theta w \rangle$$

has a maximum at  $\theta = 0$  so  $f'_w(0) = \langle Ju, w \rangle = 0$ . Similarly we have that  $\langle Jv, w \rangle = 0$ , and thus  $w \in \text{Span}\{u, v, Ju, Jv\}^\perp$ .

We then have two cases. If  $\theta_1 = 0$  then  $v = Ju$  so we can set  $u = e_1, v = Je_1$  and see that  $P = \text{Span}\{e_1, Je_1\} \times Q$  where  $Q$  is a  $2(k-1)$ -plane in  $\mathbb{C}^{n-1} = \text{Span}\{e_1, Je_1\}^\perp$ . If  $\theta_1 \neq 0$  we have that  $v = \cos \theta_1 Ju + \sin \theta_1 w$  where  $w$  is a unit vector orthogonal to  $u$  and  $Ju$ , so we can let  $u = e_1, w = e_2$  and see that  $P = \text{Span}\{e_1, \cos \theta_1 Je_1 + \sin \theta_1 e_2\} \times Q$  where  $Q$  is a  $2(k-1)$ -plane in  $\mathbb{C}^{n-2} = \text{Span}\{e_1, Je_1, e_2, Je_2\}^\perp$ .

Proceeding by induction we see that we have an oriented basis  $\{e_1, Je_1, \dots, e_n, Je_n\}$  for  $\mathbb{C}^n$  so that

$$P = \text{Span}\{e_1, \cos \theta_1 Je_1 + \sin \theta_1 e_2, \dots, e_{2k-1}, \cos \theta_k Je_{2k-1} + \sin \theta_k e_{2k}\},$$

where  $0 \leq \theta_1 \leq \dots \leq \theta_{k-1} \leq \frac{\pi}{2}$  and  $\theta_{k-1} \leq \theta_k \leq \pi - \theta_{k-1}$ .

Since we can write  $\omega = \sum_{j=1}^n e^j \wedge Je^j$  we see that  $\frac{\omega^k}{k!}$  restricts to  $P$  to give a product of  $\cos \theta_j$  which is certainly less than or equal to 1. Moreover, equality holds if and only if all of the  $\theta_j = 0$  which means that  $P$  is complex.  $\square$

Putting together Theorem 3.1 and Theorem 2.3 yields the following.

**Corollary 3.4.** *Compact complex submanifolds of Kähler manifolds are homologically volume-minimizing.*

We know that complex submanifolds are defined by holomorphic functions; i.e. solutions to the Cauchy–Riemann equations, which are a first-order PDE system, as one would expect for calibrated submanifolds.

**Example.**  $N = \{(z, \frac{1}{z}) \in \mathbb{C}^2 : z \in \mathbb{C} \setminus \{0\}\}$  is a complex curve in  $\mathbb{C}^2$ , and thus is calibrated.

**Example.** An important non-trivial example of a Kähler manifold is  $\mathbb{C}\mathbb{P}^n$ , where the zero set of a system of polynomial equations defines a (possibly singular) complex submanifold.

## 4 Special Lagrangians

Complex submanifolds are very familiar, but can we find any other interesting classes of calibrated submanifolds? The answer is that indeed we can, particularly when the manifold has special holonomy. We begin with the case of holonomy  $SU(n)$  – so-called *Calabi–Yau manifolds*. The model example for Calabi–Yau manifolds is  $\mathbb{C}^n$  with complex structure  $J$ , Kähler form  $\omega$  and holomorphic volume form

$$\Upsilon = dz_1 \wedge \dots \wedge dz_n,$$

if  $z_1, \dots, z_n$  are complex coordinates on  $\mathbb{C}^n$ .

**Theorem 4.1.** *Let  $M$  be a Calabi–Yau manifold with holomorphic volume form  $\Upsilon$ . Then  $\text{Re}(e^{-i\theta}\Upsilon)$  is a calibration for any  $\theta \in \mathbb{R}$ .*

Since  $d\Upsilon = 0$ , the result follows immediately from the following.

**Theorem 4.2.** *On  $\mathbb{C}^n$ ,  $|\Upsilon(e_1, \dots, e_n)| \leq 1$  for all unit vectors  $e_1, \dots, e_n$  with equality if and only if  $P = \text{Span}\{e_1, \dots, e_n\}$  is a Lagrangian plane, i.e.  $P$  is an  $n$ -plane such that  $\omega|_P \equiv 0$ .*

*Proof.* Let  $e_1, \dots, e_n$  be the standard basis for  $\mathbb{R}^n$  and let  $P$  be an  $n$ -plane in  $\mathbb{C}^n$ . There exists  $A \in \mathrm{GL}(n, \mathbb{C})$  so that  $f_1 = Ae_1, \dots, f_n = Ae_n$  is an orthonormal basis for  $P$ . Then  $\Upsilon(Ae_1, \dots, Ae_n) = \det_{\mathbb{C}}(A)$  so

$$|\Upsilon(f_1, \dots, f_n)|^2 = |\det_{\mathbb{C}}(A)|^2 = |\det_{\mathbb{R}}(A)| = |f_1 \wedge Jf_1 \wedge \dots \wedge f_n \wedge Jf_n| \leq |f_1| |Jf_1| \dots |f_n| |Jf_n| = 1$$

with equality if and only if  $f_1, Jf_1, \dots, f_n, Jf_n$  are orthonormal. However, this is exactly equivalent to the Lagrangian condition, since  $\omega(u, v) = g(Ju, v)$  so  $\omega|_P \equiv 0$  if and only if  $JP = P^\perp$ .  $\square$

**Definition 4.3.** A submanifold  $N$  of  $M$  calibrated by  $\mathrm{Re}(e^{-i\theta}\Upsilon)$  is called *special Lagrangian* with phase  $e^{i\theta}$ . If  $\theta = 0$  we say that  $N$  is simply special Lagrangian. By Theorem 4.2, we see that  $N$  is special Lagrangian if and only if  $\omega|_N \equiv 0$  (i.e.  $N$  is Lagrangian) and  $\mathrm{Im}\Upsilon|_N \equiv 0$  (up to a choice of orientation so that  $\mathrm{Re}\Upsilon|_N > 0$ ).

**Example.** Consider  $\mathbb{C} = \mathbb{R}^2$  with coordinates  $z = x + iy$ , complex structure  $J$  given by  $Jw = iw$ , Kähler form  $\omega = dx \wedge dy = \frac{i}{2}dz \wedge d\bar{z}$  and holomorphic volume form  $\Upsilon = dz = dx + idy$ . We want to consider the special Lagrangians in  $\mathbb{C}$ , which are 1-dimensional submanifolds or curves  $N$  in  $\mathbb{C} = \mathbb{R}^2$ .

Since  $\omega$  is a 2-form, it vanishes on any curve in  $\mathbb{C}$ . Hence every curve in  $\mathbb{C}$  is Lagrangian. For  $N$  to be special Lagrangian with phase  $e^{i\theta}$  we need that

$$\mathrm{Re}(e^{-i\theta}\Upsilon) = \cos\theta dx + \sin\theta dy$$

is the volume form on  $N$ , or equivalently that

$$\mathrm{Im}(e^{-i\theta}\Upsilon) = \cos\theta dy - \sin\theta dx$$

vanishes on  $N$ . This means that  $\cos\theta\partial_x + \sin\theta\partial_y$  is everywhere a unit tangent vector to  $N$ , so  $N$  is a straight line given by  $N = \{(t \cos\theta, t \sin\theta) \in \mathbb{R}^2 : t \in \mathbb{R}\}$  (up to translation), so it makes an angle  $\theta$  with the  $x$ -axis, hence motivating the term ‘‘phase  $e^{i\theta}$ ’’.

Notice that this result is compatible with the fact that special Lagrangians are minimal, and hence must be geodesics in  $\mathbb{R}^2$ ; i.e. straight lines.

**Example.** Consider  $\mathbb{C}^2 = \mathbb{R}^4$ . We know that  $\omega = dx_1 \wedge dy_1 + dx_2 \wedge dy_2$ . Since  $\Upsilon = dz_1 \wedge dz_2 = (dx_1 + idy_1) \wedge (dx_2 + idy_2)$ , we also know that  $\mathrm{Re}\Upsilon = dx_1 \wedge dx_2 + dy_2 \wedge dy_1$ , which looks somewhat similar. In fact, if we let  $J'$  denote the complex structure given by  $J'(\partial_{x_1}) = \partial_{x_2}$  and  $J'(\partial_{y_2}) = \partial_{y_1}$ , then  $\mathrm{Re}\Upsilon = \omega'$ , the Kähler form corresponding the complex structure  $J'$ . Hence special Lagrangians in  $\mathbb{C}^2$  are complex curves for a different complex structure.

In fact, we have a hyperkähler triple of complex structures  $J_1, J_2, J_3$ , where  $J_1 = J$  is the standard one and  $J_3 = J_1J_2 = -J_2J_1$  so that  $J_1 = J_2J_3 = -J_3J_2$  and  $J_2 = J_3J_1 = -J_1J_3$ , and the corresponding Kähler forms are  $\omega = \omega_1, \omega_2, \omega_3$  which are orthogonal and the same length with  $\Upsilon = \omega_2 + i\omega_3$ .

This shows we should only consider complex dimension 3 and higher to find new calibrated submanifolds.

**Example.** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a smooth function and let  $N = \mathrm{Graph}(f) \subseteq \mathbb{R}^{2n} = \mathbb{C}^n$ . We want to see when  $N$  is special Lagrangian. We see that tangent vectors to  $N$  are given by

$$e_1 + i\nabla_{e_1}f, \dots, e_n + i\nabla_{e_n}f.$$

Hence  $N$  is Lagrangian if and only if

$$\omega(e_j + i\nabla_{e_j}f, e_k + i\nabla_{e_k}f) = \nabla_{e_k}f_j - \nabla_{e_j}f_k = 0$$

for all  $j, k$ . Since  $\mathbb{R}^n$  is simply connected, this occurs if and only if there exists  $F$  such that  $f_j = \nabla_{e_j}F$ ; i.e.  $f = \nabla F$ .

Recall that  $\Upsilon = dz_1 \wedge \dots \wedge dz_n$ . We know that  $N$  is special Lagrangian if and only if  $N$  is Lagrangian and  $\mathrm{Im}\Upsilon$  vanishes on  $N$ . Now

$$\Upsilon(a_1 + ib_1, \dots, a_n + ib_n) = \det_{\mathbb{C}}(A + iB)$$

where  $A, B$  are the matrices with columns  $a_i, b_j$  respectively. Hence

$$\Upsilon(e_1 + i\nabla_{e_1}\nabla F, \dots, e_n + i\nabla_{e_n}\nabla F) = \det_{\mathbb{C}}(I + i\text{Hess } F),$$

where  $\text{Hess } F = (\frac{\partial^2 F}{\partial x_i \partial x_j})$ .

Therefore  $N = \text{Graph}(f)$  is special Lagrangian (up to a choice of orientation) if and only if  $f = \nabla F$  and

$$\text{Im } \det_{\mathbb{C}}(I + i\text{Hess } F) = 0.$$

If  $n = 2$ ,

$$I + i\text{Hess } F = \begin{pmatrix} 1 + iF_{xx} & iF_{xy} \\ iF_{yx} & 1 + iF_{yy} \end{pmatrix}.$$

Therefore, the determinant gives

$$1 - F_{xx}F_{yy} + F_{xy}^2 + i(F_{xx} + F_{yy}),$$

then the imaginary part is  $F_{xx} + F_{yy}$ . Therefore,  $N$  is special Lagrangian if and only if  $\Delta F = 0$ .

As we know, a graph in  $\mathbb{C}^2$  of  $f = u + iv : \mathbb{C} \rightarrow \mathbb{C}$  is a complex surface if and only if  $u + iv$  is holomorphic, which implies that  $u, v$  are harmonic. We know that special Lagrangians in  $\mathbb{C}^2$  are complex surfaces for a different complex structure, so this is expected.

If  $n = 3$ ,

$$I + i\text{Hess } F = \begin{pmatrix} 1 + iF_{xx} & iF_{xy} & iF_{xz} \\ iF_{yx} & 1 + iF_{yy} & iF_{yz} \\ iF_{zx} & iF_{zy} & 1 + iF_{zz} \end{pmatrix}.$$

Hence,

$$\begin{aligned} \text{Im } \det_{\mathbb{C}}(I + i\text{Hess } F) &= F_{xx} + F_{yy} + F_{zz} \\ &\quad - F_{xx}(F_{yy}F_{zz} - F_{yz}^2) - F_{xy}(F_{yz}F_{zx} - F_{xy}F_{zz}) - F_{zx}(F_{xy}F_{yz} - F_{yy}F_{zx}). \end{aligned}$$

Therefore,  $N$  is special Lagrangian if and only if

$$\begin{aligned} -\Delta F &= F_{xx} + F_{yy} + F_{zz} \\ &= F_{xx}(F_{yy}F_{zz} - F_{yz}^2) - F_{xy}(F_{xy}F_{zz} - F_{yz}F_{zx}) + F_{zx}(F_{xy}F_{yz} - F_{yy}F_{zx}) \\ &= \det \text{Hess } F. \end{aligned}$$

We now wish to describe some very important examples of special Lagrangians, which are asymptotic to pairs of planes.

**Example.**  $\text{SU}(n)$  acts transitively on the space of special Lagrangian planes with isotropy  $\text{SO}(n)$ . So any special Lagrangian plane is given by  $A \cdot \mathbb{R}^n$  for  $A \in \text{SU}(n)$  where  $\mathbb{R}^n$  is the standard real  $\mathbb{R}^n$  in  $\mathbb{C}^n$ .

Given  $\theta = (\theta_1, \dots, \theta_n)$  we can define a plane  $P(\theta) = \{(e^{i\theta_1}x_1, \dots, e^{i\theta_n}x_n) \in \mathbb{C}^n : (x_1, \dots, x_n) \in \mathbb{R}^n\}$  (where we can swap orientation). We see that  $P(\theta)$  is special Lagrangian if and only if  $\text{Re } \Upsilon|_P = \pm \cos(\theta_1 + \dots + \theta_n) = 1$  so that  $\theta_1 + \dots + \theta_n \in \pi\mathbb{Z}$ . Given any  $\theta_1, \dots, \theta_n \in (0, \pi)$  with  $\theta_1 + \dots + \theta_n = \pi$ , there exists a special Lagrangian  $N$  (called a *Lawlor neck*) asymptotic to  $P(0) \cup P(\theta)$ : see, for example, [42, Example 8.3.15] or §8 for details. It is diffeomorphic to  $\mathcal{S}^{n-1} \times \mathbb{R}$ . By rotating coordinates we have a special Lagrangian with phase  $i$  asymptotic to  $P(-\frac{\theta}{2}) \cup P(\frac{\theta}{2})$ .

The simplest case is when  $\theta_1 = \dots = \theta_n = \frac{\pi}{n}$ : here  $N$  is called the *Lagrangian catenoid*. When  $n = 2$ , under a coordinate change the Lagrangian catenoid becomes the complex curve  $\{(z, \frac{1}{z}) \in \mathbb{C}^2 : z \in \mathbb{C} \setminus \{0\}\}$  that we saw before. When  $n = 3$ , the only possibilities for the angles are  $\sum_i \theta_i = \pi, 2\pi$ , but if  $\sum_i \theta_i = 2\pi$  we can rotate coordinates and change the order of the planes so that  $P(0) \cup P(\theta)$  becomes  $P(0) \cup P(\theta')$  where  $\sum_i \theta'_i = \pi$ . Hence, given any pair of transverse special Lagrangian planes in  $\mathbb{C}^3$ , there exists a Lawlor neck asymptotic to their union.

**Remark.** Using complex geometry it is easy to classify all of the smooth special Lagrangians in  $\mathbb{C}^2$  asymptotic to a pair of transverse planes, and one sees that the Lawlor necks in  $\mathbb{C}^2$  are the unique exact special Lagrangians with this property. It is now known that the Lawlor necks are the unique smooth exact special Lagrangian asymptotic to a pair of planes in all dimensions [25].



We can find special Lagrangians in Calabi–Yau manifolds using the following easy result.

**Proposition 4.4.** *Let  $(M, \omega, \Upsilon)$  be a Calabi–Yau manifold and let  $\sigma : M \rightarrow M$  be such that  $\sigma^2 = \text{Id}$ ,  $\sigma^*(\omega) = -\omega$ ,  $\sigma^*(\Upsilon) = \bar{\Upsilon}$ . Then  $\text{Fix}(\sigma)$  is special Lagrangian, if it is non-empty.*

**Example.** Let  $X = \{[z_0, \dots, z_4] \in \mathbb{C}\mathbb{P}^4 : z_0^5 + \dots + z_4^5 = 0\}$  (the *Fermat quintic*) with its Calabi–Yau structure (which exists by Yau’s solution of the Calabi conjecture since the first Chern class of  $X$  vanishes). Let  $\sigma$  be the restriction of complex conjugation on  $\mathbb{C}\mathbb{P}^4$  to  $X$ . Then the fixed point set of  $\sigma$ , which is the real locus in  $X$ , is a special Lagrangian 3-fold (if it is non-empty). (There is a subtlety here:  $\sigma$  is certainly an anti-holomorphic isometric involution for the induced metric on  $X$ , but this is *not* the same as the Calabi–Yau metric on  $X$ . Nevertheless, it is the case that  $\sigma$  satisfies the conditions of Proposition 4.4.)

**Example.** There exists a Calabi–Yau metric on  $T^*\mathcal{S}^n$  (the Stenzel metric [78]) so that the base  $\mathcal{S}^n$  is special Lagrangian: When  $n = 2$  this is a hyperkähler metric called the Eguchi–Hanson metric [11].

## 5 Constructing calibrated submanifolds

It is easy to construct complex submanifolds in Kähler manifolds algebraically. Constructing other calibrated submanifolds is much more challenging because one needs to solve a nonlinear PDE, even in Euclidean space. There are approaches in Euclidean space and other simple spaces which have involved reducing the problem to ODEs or other problems which do not require PDE (for example, algebraic methods). For example, we have the following methods, which you can find out more about in [42] or the references provided.

- Symmetries/evolution equations [17, 20, 21, 28, 30, 31, 33, 34, 39, 40, 41, 52, 54].
- Use of integrable systems to study calibrated cones [8, 9, 22, 43, 65].
- Calibrated cones and ruled smoothings of these cones [2, 4, 13, 14, 32, 52, 53, 59].
- Vector sub-bundle constructions [27, 45, 46].
- Classification of calibrated submanifolds satisfying pointwise constraints on their second fundamental form [4, 12, 26, 59, 60].

However, an important direction which has borne fruit in calibrated geometry and special holonomy recently has been to study the nonlinear PDE head on, especially by perturbative and gluing methods.

We want to solve nonlinear PDE, so how do we tackle this? The idea is to use the linear case to help. Suppose we are on a compact manifold  $N$  and recall the theory of linear *elliptic* operators  $L$  of order  $l$  on  $N$ , including:

- the definition of ellipticity of  $L$  via the *principal symbol*  $\sigma_L$  (which encodes the highest order derivatives in the operator) being an isomorphism;
- the use of *Hölder spaces*  $C^{k,a}$  to give elliptic regularity theory (so-called *Schauder theory*), namely that if  $w \in C^{k,a}$  and  $Lv = w$  then  $v \in C^{k+l,a}$  and there is a universal constant  $C$  so that

$$\|v\|_{C^{k+l,a}} \leq C(\|Lv\|_{C^{k,a}} + \|v\|_{C^0})$$

(and we can drop the  $\|v\|_{C^0}$  term if  $v$  is orthogonal to  $\text{Ker } L$ );

- the adjoint operator  $L^*$  and that  $\sigma_{L^*} = (-1)^l \sigma_L^*$  so that  $L^*$  is elliptic if and only if  $L$  is elliptic; and
- the Fredholm theory of  $L$ , namely that  $\text{Ker } L$  (and hence  $\text{Ker } L^*$ ) is finite-dimensional, and we can solve  $Lv = w$  if and only if  $w \in (\text{Ker } L^*)^\perp$ .

We shall discuss this in a model example which we shall use throughout this section.

**Example.** The Laplacian on functions is given by  $\Delta f = d^*df$  which in normal coordinates at a point is given by  $f \mapsto -\sum_i \frac{\partial^2 f}{\partial x_i^2}$ , so it is a linear second order differential operator. We see that its principal symbol is  $\sigma_\Delta(x, \xi)f = -|\xi|^2 f$  which is an isomorphism for  $\xi \in T_x^*N \setminus \{0\}$ , so  $\Delta$  is elliptic. We therefore have that if  $h \in C^{k,a}(N)$  and  $\Delta f = h$  then  $f \in C^{k+2,a}(N)$ , and we have an estimate

$$\|f\|_{C^{k+2,a}} \leq C(\|\Delta f\|_{C^{k,a}} + \|f\|_{C^0}).$$

We also know that  $\Delta^* = \Delta$  and  $\text{Ker } \Delta$  is given by the constant functions (since if  $f \in \text{Ker } \Delta$  then

$$0 = \langle f, \Delta f \rangle_{L^2} = \langle f, d^*df \rangle_{L^2} = \|df\|_{L^2}^2$$

so  $df = 0$ ). Hence, we can solve  $\Delta f = h$  if and only if  $h$  is orthogonal to the constants, i.e.  $\int_N h \text{vol}_N = 0$ .

The operator defining the minimal graph equation for a hypersurface is

$$P(f) = -\text{div} \left( \frac{\nabla f}{\sqrt{1 + |\nabla f|^2}} \right),$$

which is a nonlinear second order operator whose linearisation  $L_0P$  at 0 is  $\Delta$ . Thus  $P$  is a nonlinear elliptic operator at 0. If we linearise  $P$  at  $f_0$  we find a more complicated expression depending on  $f_0$ , but it is still a perturbation of the Laplacian.

Suppose we are on a compact manifold  $N$  and we want to solve  $P(f) = 0$  where  $P$  is the minimal graph operator on functions  $f$ . Let us consider regularity for  $f$ . We can re-arrange  $P(f) = 0$  by taking all of the second derivatives to one side as:

$$R(x, \nabla f(x)) \nabla^2 f(x) = E(x, \nabla f(x))$$

where  $x \in N$ . Since  $L_0P = \Delta$  is elliptic and ellipticity is an open condition we know that the operator  $L_f$  (depending on  $f$ ) given by

$$L_f(h)(x) = R(x, \nabla f(x)) \nabla^2 h(x)$$

is a *linear* elliptic operator whenever  $\|\nabla f\|_{C^0}$  is small, in particular if  $\|f\|_{C^{1,a}}$  is sufficiently small. The operator  $L_f$  does not have smooth coefficients, but if  $f \in C^{k,a}$  then the coefficients  $R \in C^{k-1,a}$ .

Suppose that  $f \in C^{1,a}$  and  $\|f\|_{C^{1,a}}$  is small with  $P(f) = 0$ . Then  $L_f(f) = E(f)$  and  $L_f$  is a linear *second order* elliptic operator with coefficients in  $C^{0,a}$  and  $E(f)$  is in  $C^{0,a}$ . So by elliptic regularity we can deduce that  $f \in C^{2,a}$ . We have gained one degree of regularity, so we can “bootstrap”, i.e. proceed by induction and deduce that any  $C^{1,a}$  solution to  $P(f) = 0$  is smooth.

**Example.**  $C^{1,a}$ -minimal submanifolds (and thus calibrated submanifolds) are *smooth*.

**Remark.** More sophisticated techniques can be used to deduce that  $C^1$ -minimal submanifolds are real analytic [69]. Notice that elliptic regularity results are *not* valid for  $C^k$  spaces, so this result is not obvious.

We can also arrange our simple equation  $P(f) = 0$  as  $\Delta f + Q(\nabla f, \nabla^2 f) = 0$ , where  $Q$  is nonlinear but linear in  $\nabla^2 f$ . If we know that  $\int_N P(f) \text{vol}_N = 0$ , i.e. that  $P(f)$  is orthogonal to the constants, then we can always solve  $\Delta f_0 = -Q(\nabla f, \nabla^2 f)$ . We do know that  $\int_N P(f) \text{vol}_N = 0$  since  $P$  has a divergence form. This means we are in the setting for implementing the Implicit Function Theorem for Banach spaces to conclude that we can always solve  $P(f) = 0$  for some  $f$  near 0, and  $f$  will be smooth by our regularity argument above. In general, we will use the following.

**Theorem 5.1** (Implicit Function Theorem). *Let  $X, Y$  be Banach spaces, let  $U \ni 0$  be open in  $X$ , let  $P : U \rightarrow Y$  with  $P(0) = 0$  and  $L_0P : X \rightarrow Y$  surjective with finite-dimensional kernel  $K$ .*

*Then for some  $U$ ,  $P^{-1}(0) = \{u \in U : P(u) = 0\}$  is a manifold of dimension  $\dim K$ . Moreover, if we write  $X = K \oplus Z$ ,  $P^{-1}(0) = \text{Graph } G$  for some map  $G$  from an open set in  $K$  to  $Z$  with  $G(0) = 0$ .*

This gives us a way to describe all perturbations of a given calibrated submanifold, as we now see in the special Lagrangian case, due to McLean [66].

**Theorem 5.2.** *Let  $N$  be a compact special Lagrangian in a Calabi–Yau manifold  $M$ . Then the moduli space of deformations of  $N$  is a smooth manifold of dimension  $b^1(N)$ .*

**Remark.** One should compare this result to the deformation theory for complex submanifolds in Kähler manifolds. There, one does not get that the moduli space is a smooth manifold: in fact, it can be singular, and one has *obstructions* to deformations. It is somewhat remarkable that special Lagrangian calibrated geometry enjoys a much better deformation theory than this classical calibrated geometry. The deformation theory of embedded compact complex submanifolds in Calabi–Yau manifolds has recently been revisited using analytic techniques [67].

*Proof.* The tubular neighbourhood theorem gives us a diffeomorphism  $\exp : S \subseteq \nu(N) \rightarrow T \subseteq M$  which maps the zero section to  $N$ ; in other words, we can write any nearby submanifold to  $N$  as the graph of a normal vector field on  $N$ . We know that  $N$  is Lagrangian, so the complex structure  $J$  gives an isomorphism between  $\nu(N)$  and  $TN$  and the metric gives an isomorphism between  $TN$  and  $T^*N$ :  $v \mapsto g(Jv, \cdot) = \omega(v, \cdot) = \alpha_v$ . Therefore any deformation of  $N$  in  $T$  is given as the graph of a 1-form. In fact, using the Lagrangian neighbourhood theorem, we can arrange that any  $N' \in T$  is the graph of a 1-form  $\alpha$ , so that if  $f_\alpha : N \rightarrow N_\alpha$  is the natural diffeomorphism then

$$f_\alpha^*(\omega) = d\alpha \quad \text{and} \quad - * f_\alpha^*(\text{Im } \Upsilon) = F(\alpha, \nabla \alpha) = d^* \alpha + Q(\alpha, \nabla \alpha),$$

where the second formula follows from a calculation using the special Lagrangian condition on  $N$  and the fact that the ambient structure is Calabi–Yau. Hence,  $N_\alpha$  is special Lagrangian if and only if  $P(\alpha) = (F(\alpha, \nabla \alpha), d\alpha) = 0$ . This means that infinitesimal special Lagrangian deformations are given by closed and coclosed 1-forms, which is the kernel of  $L_0P$ .

Since  $\text{Im } \Upsilon = 0$  on  $N$  we have that  $[\text{Im } \Upsilon] = 0$  on  $N_\alpha$ , which means that  $f_\alpha^*(\text{Im } \Upsilon)$  is exact. Thus  $F(\alpha, \nabla \alpha) = - * f_\alpha^*(\text{Im } \Upsilon)$  is coexact and so

$$P : C^\infty(S) \rightarrow d^*(C^\infty(T^*N)) \oplus d(C^\infty(T^*N)) \subseteq C^\infty(\Lambda^0 T^*N \oplus \Lambda^2 T^*N).$$

If we let  $X = C^{1,a}(T^*N)$ ,  $Y = d^*(C^{1,a}(T^*N)) \oplus d(C^{1,a}(T^*N))$  and  $U = C^{1,a}(S)$  we can apply the Implicit Function Theorem if we know that

$$L_0P : \alpha \in X \mapsto (d^* \alpha, d\alpha) \in Y$$

is surjective, i.e. given  $d\beta + d^* \gamma \in Y$  does there exist  $\alpha$  such that  $d\alpha = d\beta$  and  $d^* \alpha = d^* \gamma$ ? If we let  $\alpha = \beta + df$  then we need  $\Delta f = d^* df = d^*(\gamma - \beta)$ . Since

$$\int_N d^*(\gamma - \beta) \text{vol}_N = \pm \int_N d * (\gamma - \beta) = 0$$

we can solve the equation for  $f$ , and hence  $L_0P$  is surjective.

Therefore  $P^{-1}(0)$  is a manifold of dimension  $\dim \text{Ker } L_0P = b^1(N)$  by Hodge theory. Moreover, if  $P(\alpha) = 0$  then  $N_\alpha$  is special Lagrangian, hence minimal and since  $\alpha \in C^{1,a}$  we deduce that  $\alpha$  is in fact smooth.  $\square$

**Example.** The special Lagrangian  $S^n$  in  $T^*S^n$  has  $b^1 = 0$  and so is rigid.

Observe that if we have a special Lagrangian  $T^n$  in  $M$  then  $b^1(T^n) = n$  and, if the torus is close to flat then its deformations locally foliate  $M$  (as there will be  $n$  nowhere vanishing harmonic 1-forms), so we can hope to find special Lagrangian torus fibrations. This cannot happen in compact manifolds without singular fibres, but still motivates the SYZ conjecture in Mirror Symmetry. The deformation result also motivates the following theorem [3].

**Theorem 5.3.** *Every compact oriented real analytic Riemannian 3-manifold can be isometrically embedded in a Calabi–Yau 3-fold as the fixed point set of an involution.*

**Remark.** Theorem 5.2 has also been extended to certain non-compact, singular and boundary settings, for example in [6, 36, 72].

Another well-known way to get a solution of a linear PDE from two solutions is simply to add them. However, for a nonlinear PDE  $P(v) = 0$  this will not work. Intuitively, we can try to add two solutions to give us a solution  $v_0$  for which  $P(v_0)$  is small. Then we may try to perturb  $v_0$  by  $v$  to solve  $P(v + v_0) = 0$ .

Geometrically, this occurs when we have two calibrated submanifolds  $N_1, N_2$  and then glue them together to give a submanifold  $N$  which is “almost” calibrated, then we deform  $N$  to become calibrated. If the two submanifolds  $N_1, N_2$  are glued using a very long neck then one can imagine that  $N$  is almost the disjoint union of  $N_1, N_2$  and so close to being calibrated. If instead one scales  $N_2$  by a factor  $t$  and then glues it into a singular point of  $N_1$ , we can again imagine that as  $t$  becomes very small  $N$  resembles  $N_1$  and so again is close to being calibrated. These two examples are in fact related, because if we rescale the shrinking  $N_2$  to fixed size, then we get a long neck between  $N_1$  and  $N_2$  of length of order  $-\log t$ . However, although these pictures are appealing, they also reveal the difficulty in this approach: as  $t$  becomes small,  $N$  becomes more “degenerate”, giving rise to analytic difficulties which are encoded in the geometry of  $N_1, N_2$  and  $N$ .

These ideas are used extensively in geometry, and particularly successfully in calibrated geometry e.g. [7, 23, 35, 37, 38, 51, 57, 62, 73]. A particular simple case is the following, which we will describe to show the basic idea of the gluing method.

**Theorem 5.4.** *Let  $N$  be a compact connected 3-manifold and let  $i : N \rightarrow M$  be a special Lagrangian immersion with transverse self-intersection points in a Calabi–Yau manifold  $M$ . Then there exist embedded special Lagrangians  $N_t$  such that  $N_t \rightarrow N$  as  $t \rightarrow 0$ .*

**Remark.** One might ask about the sense of convergence here: for definiteness, we can say that  $N_t$  converges to  $N$  in the sense of currents; that is, if we have any compactly supported 3-form  $\chi$  on  $M$  then  $\int_{N_t} \chi \rightarrow \int_N \chi$  as  $t \rightarrow 0$ . However, all sensible notions of convergence of submanifolds will be true in this setting.

*Proof.* Here we only provide a sketch of the proof: see, for example, [35, §9] for a detailed proof.

At each self-intersection point of  $N$  the tangent spaces are a pair of transverse 3-planes, which we can view as a pair of transverse special Lagrangian 3-planes  $P_1, P_2$  in  $\mathbb{C}^3$ . Since we are in dimension 3, we know that there exists a (unique up to scale) special Lagrangian Lawlor neck  $L$  asymptotic to  $P_1 \cup P_2$ . We can then glue  $tL$  into  $N$  near each intersection point to get a compact embedded submanifold  $S_t = N \# tL$  (if we glue in a Lawlor neck for every self-intersection point). We can also arrange that  $S_t$  is Lagrangian, i.e. that it is a Lagrangian connect sum.

Now we want to perturb  $S_t$  to be special Lagrangian. Since  $S_t$  is Lagrangian, by the deformation theory we can write any nearby submanifold as the graph of a 1-form  $\alpha$ , and this graph will be special Lagrangian if and only if (using the same notation as in our deformation theory discussion)

$$P_t(\alpha) = (- * f_\alpha^*(\text{Im } \Upsilon), f_\alpha^*(\omega)) = 0.$$

Since  $S_t$  is Lagrangian but not special Lagrangian we have that

$$f_\alpha^*(\omega) = d\alpha \quad \text{and} \quad - * f_\alpha^*(\text{Im } \Upsilon) = P_t(0) + d_t^* \alpha + Q_t(\alpha, \nabla \alpha)$$

where  $P_t(0) = - * \text{Im } \Upsilon|_{S_t}$  and  $d_t^* = L_0 P_t$ , which is a perturbation of the usual  $d^*$  since we are no longer linearising at a point where  $P_t(0) = 0$ . By choosing  $\alpha = df$ , we then have to solve

$$\Delta_t f = -P_t(0) - Q_t(\nabla f, \nabla^2 f)$$

where  $\Delta_t$  is a perturbation of the Laplacian.

For simplicity, let us suppose that  $\Delta_t$  is the Laplacian on  $S_t$ . The idea is to view our equation as a fixed point problem. We know that if we let  $X^k = \{f \in C^{k,a}(N) : \int_N f \text{ vol}_N = 0\}$  then  $\Delta_t : X^{k+2} \rightarrow X^k$  is an isomorphism so it has an inverse  $G_t$ . We know by our elliptic regularity result that there exists a constant  $C(\Delta_t)$  such that

$$\|f\|_{C^{k+2,a}} \leq C(\Delta_t) \|\Delta_t f\|_{C^{k,a}} \Leftrightarrow \|G_t h\|_{C^{k+2,a}} \leq C(\Delta_t) \|h\|_{C^{k,a}}$$

for any  $f \in X^{k+2}$ ,  $h \in X^k$ .

We thus see that  $P_t(f) = 0$  for  $f \in X^{k+2}$  if and only if

$$f = G_t(-P_t(0) - Q_t(f)) = F_t(f).$$

The idea is now to show that  $F_t$  is a contraction sufficiently near 0 for all  $t$  small enough. Then it will have a (unique) fixed point near 0, which will also be smooth because it satisfies  $P_t(f) = 0$  and hence defines a special Lagrangian as the graph of  $df$  over  $S_t$ .

We know that  $F_t : X^{k+2} \rightarrow X^{k+2}$  with

$$\|F_t(f_1) - F_t(f_2)\|_{C^{k+2,a}} = \|G_t(Q_t(f_1) - Q_t(f_2))\|_{C^{k+2,a}} \leq C(\Delta_t)\|Q_t(f_1) - Q_t(f_2)\|_{C^{k,a}}.$$

Since  $Q_t$  and its first derivatives vanish at 0 we know that

$$\|Q_t(f_1) - Q_t(f_2)\|_{C^{k,a}} \leq C(Q_t)\|f_1 - f_2\|_{C^{k+2,a}}(\|f_1\|_{C^{k+2,a}} + \|f_2\|_{C^{k+2,a}}).$$

We deduce that

$$\|F_t(f_1) - F_t(f_2)\|_{C^{k+2,a}} \leq C(\Delta_t)C(Q_t)\|f_1 - f_2\|_{C^{k+2,a}}(\|f_1\|_{C^{k+2,a}} + \|f_2\|_{C^{k+2,a}})$$

and

$$\|F_t(0)\|_{C^{k+2,a}} = \|G_t(P_t(0))\|_{C^{k+2,a}} \leq C(\Delta_t)\|P_t(0)\|_{C^{k,a}}.$$

Hence,  $F_t$  is a contraction on  $\overline{B_{\epsilon_t}(0)} \subseteq X^{k+2}$  if we can choose  $\epsilon_t$  so that

$$2C(\Delta_t)\|P_t(0)\|_{C^{k,a}} \leq \epsilon_t \leq \frac{1}{4C(\Delta_t)C(Q_t)}.$$

(This also proves Theorem 5.2, where we used the Implicit Function Theorem, by hand since there  $P_t(0) = P(0) = 0$  so we just need to take  $\epsilon_t$  small enough.) In other words, we need that

- $P_t(0)$  is small, so  $S_t$  is “close” to being calibrated and is a good approximation to  $P_t(f) = 0$ ;
- $C(\Delta_t), C(Q_t)$ , which are determined by the linear PDE and geometry of  $N, L$  and  $S_t$ , are well-controlled as  $t \rightarrow 0$ .

The statement of the theorem is then that there exists  $t$  sufficiently small and  $\epsilon_t$  so that the contraction mapping argument works.

This is a delicate balancing act since as  $t \rightarrow 0$  parts of the manifold are collapsing, so the constants  $C(\Delta_t), C(Q_t)$  above (which depend on  $t$ ) can and typically do blow-up as  $t \rightarrow 0$ . To control this, we need to understand the Laplacian on  $N, L$  and  $S_t$  and introduce “weighted” Banach spaces so that  $tL$  gets rescaled to constant size (independent of  $t$ ), and  $S_t$  resembles the union of two manifolds with a cylindrical neck (as we described earlier). It is also crucial to understand the relationship between the kernels and cokernels of the Laplacian on the *non-compact*  $N$  (with the intersection points removed),  $L$  and compact  $S_t$ : here is where connectedness is important so that the kernel and cokernel of the Laplacian is 1-dimensional.  $\square$

**Remark.** In more challenging gluing problems it is not possible to show that the relevant map is a contraction, but rather one can instead appeal to an alternative theorem (e.g. Schauder fixed point theorem) to show that it still has a fixed point.

## 6 Associative and coassociative submanifolds

We now want to introduce our calibrated geometry associated with  $G_2$  holonomy. The first key result is the following.

**Theorem 6.1.** *Let  $(M^7, \varphi)$  be a  $G_2$  manifold (so  $\varphi$  is a closed and coclosed positive 3-form). Then  $\varphi$  and  $*\varphi$  are calibrations.*

*Proof.* Let  $u, v, w$  be oriented orthonormal vectors in  $\mathbb{R}^7$ . There exists an element  $A$  of  $G_2$  so that  $Au = e_1$ . The subgroup of  $G_2$  fixing  $e_1$  is isomorphic to  $SU(3)$ , and we know from the proof of Wirtinger's inequality (Theorem 3.2) there exists a (special) unitary transformation so that  $v = e_2$  and  $w = \cos \theta e_3 + \sin \theta v$  for some  $\theta$  and  $v$  orthogonal to  $e_1, e_2, e_3$ . Since  $\varphi(e_1, e_2, \cdot) = dx_3$ , we see that  $\varphi(u, v, w) = \cos \theta$ . Hence, since  $\varphi$  is closed,  $\varphi$  is a calibration and the calibrated planes are given by  $A \cdot \text{Span}\{e_1, e_2, e_3\}$  for  $A \in G_2$ .

By Lemma 3.3,  $*\varphi$  is also a calibration.  $\square$

Let us look at the calibrated planes and start with  $\varphi$ , which we take to be the following on  $\mathbb{R}^7$ :

$$\varphi = dx_{123} + dx_{145} + dx_{167} + dx_{246} - dx_{257} - dx_{347} - dx_{356},$$

where we use the short-hand notation  $dx_{i_1 \dots i_k} = dx_{i_1} \wedge dx_{i_2} \wedge \dots \wedge dx_{i_k}$ . Hence,  $*\varphi$  on  $\mathbb{R}^7$  is given by:

$$*\varphi = dx_{4567} + dx_{2367} + dx_{2345} + dx_{1357} - dx_{1346} - dx_{1256} - dx_{1247}.$$

If  $u, v, w$  are unit vectors in  $\mathbb{R}^7 \cong \text{Im } \mathbb{O}$  (the imaginary octonions), then  $\varphi(u, v, w) = \langle u \times v, w \rangle = 1$  if and only if  $w = u \times v$ , so  $P = \text{Span}\{u, v, w\}$  is a copy of  $\text{Im } \mathbb{H}$  in  $\text{Im } \mathbb{O}$ ; in other words,  $\text{Span}\{1, u, v, w\}$  is an associative subalgebra of  $\mathbb{O}$ . Moreover, suppose we define a vector-valued 3-form  $\chi$  on  $\mathbb{R}^7$  by

$$\chi(u, v, w) = [u, v, w] = u(vw) - (uv)w,$$

known as the *associator*. Then we observe the following.

**Lemma 6.2.** *A 3-plane  $P$  in  $\mathbb{R}^7$  satisfies  $\chi|_P \equiv 0$  if and only if  $P$  admits an orientation so that it is calibrated by  $\varphi$ .*

*Proof.* Since the associator is clearly invariant under  $G_2$  we can put any plane  $P$  in standard position using  $G_2$ , i.e. as in the proof of Theorem 6.1, we can write  $P = \text{Span}\{e_1, e_2, \cos \theta e_3 + \sin \theta v\}$  for some  $v$  orthogonal to  $e_1, e_2, e_3$ . We can calculate that  $[e_1, e_2, e_3] = 0$  whereas  $[e_1, e_2, v] \neq 0$  for any  $v$  orthogonal to  $e_1, e_2, e_3$ . Moreover,  $P$  is calibrated by  $\varphi$  if and only if  $\theta = 0$ . We thus see that  $P$  is calibrated by  $\varphi$  (up to a choice a orientation) if and only if  $\chi|_P \equiv 0$ .  $\square$

Hence we call the  $\varphi$ -calibrated planes *associative*. In general on a  $G_2$  manifold we can define a 3-form  $\chi$  with values in  $TM$  using the pointwise formula above.

For  $*\varphi$  we see that  $*\varphi|_P = \text{vol}_P$  for a plane  $P$  if and only if  $\varphi|_{P^\perp} = \text{vol}_{P^\perp}$ . Hence the planes calibrated by  $*\varphi$  are the orthogonal complements of the associative planes, so we call them *coassociative*. We have a similar alternative characterisation for 4-planes calibrated by  $*\varphi$ .

**Lemma 6.3.** *A 4-plane  $P$  in  $\mathbb{R}^7$  satisfies  $\varphi|_P \equiv 0$  if and only if  $P$  admits an orientation so that it is calibrated by  $*\varphi$ .*

*Proof.* We know that given a 4-plane  $P$  we can choose coordinates such that  $P^\perp = \text{Span}\{e_1, e_2, \cos \theta e_3 + \sin \theta(a_4 e_4 + a_5 e_5 + a_6 e_6 + a_7 e_7)\}$  where  $\sum_j a_j^2 = 1$ . Then

$$P = \text{Span}\{-\sin \theta e_3 + \cos \theta(a_j e_j), a_5 e_4 - a_4 e_5 + a_7 e_6 - a_6 e_7, \\ a_6 e_4 - a_7 e_5 - a_4 e_6 + a_5 e_7, a_7 e_4 + a_6 e_5 - a_5 e_6 - a_4 e_7\}.$$

We can then see directly that  $*\varphi|_P = \cos \theta$ . We also have  $\varphi(e_i, e_j, e_k) = 0$  for  $i, j, k \in \{4, 5, 6, 7\}$  and  $e_3 \lrcorner \varphi = -dx_{47} - dx_{56}$ , so that  $\varphi(-\sin \theta e_3 + \cos \theta(a_j e_j), v, w)$  is a non-zero multiple of  $\sin \theta$  for some  $v, w \in P$ . Hence  $\varphi|_P = 0$  if and only if  $\theta = 0$ , which is if and only if  $P$  is calibrated by  $*\varphi$  (again up to a choice of orientation).  $\square$

We thus can define our calibrated submanifolds.

**Definition 6.4.** Submanifolds in  $(M^7, \varphi)$  calibrated by  $\varphi$  are called *associative* 3-folds. Moreover,  $N$  is associative if and only if  $\chi|_N \equiv 0$  (up to a choice of orientation).

Submanifolds in  $(M^7, \varphi)$  calibrated by  $*\varphi$  are called *coassociative* 4-folds. Moreover,  $N$  is coassociative if and only if  $\varphi|_N \equiv 0$  (up to a choice of orientation).

It is instructive to see the form that the associative or coassociative condition takes by studying associative or coassociative graphs in  $\mathbb{R}^7$ : see [20] for details.

A simple way to get associative and coassociative submanifolds is by using known geometries.

**Proposition 6.5.** *Let  $x_1, \dots, x_7$  be coordinates on  $\mathbb{R}^7$  and let  $z_j = x_{2j} + ix_{2j+1}$  be coordinates on  $\mathbb{C}^3$  so that  $\mathbb{R}^7 = \mathbb{R} \times \mathbb{C}^3$ .*

- (a)  *$N = \mathbb{R} \times S \subseteq \mathbb{R} \times \mathbb{C}^3$  is associative or coassociative if and only if  $S$  is a complex curve or a special Lagrangian 3-fold with phase  $-i$ , respectively.*
- (b)  *$N \subseteq \{0\} \times \mathbb{C}^3$  is associative or coassociative if and only if  $N$  is a special Lagrangian 3-fold or a complex surface, respectively.*

*Proof.* Recall the Kähler form  $\omega$  and holomorphic volume form  $\Upsilon$  on  $\mathbb{C}^3$ . We can write

$$\varphi = dx_1 \wedge \omega + \operatorname{Re} \Upsilon \quad \text{and} \quad * \varphi = \frac{1}{2} \omega^2 - dx_1 \wedge \operatorname{Im} \Upsilon.$$

For associatives, we see that  $\varphi|_{\mathbb{R} \times S} = dx_1 \wedge \operatorname{vol}_S$  if and only if  $\omega|_S = \operatorname{vol}_S$  and  $\varphi|_N = \operatorname{Re} \Upsilon|_N$  for  $N \subseteq \mathbb{C}^3$ . For coassociatives, we see that  $*\varphi|_{\mathbb{R} \times S} = dx_1 \wedge \operatorname{vol}_S$  if and only if  $-\operatorname{Im} \Upsilon|_S = \operatorname{vol}_S$  and  $*\varphi|_N = \frac{1}{2} \omega^2|_N$  for  $N \subseteq \mathbb{C}^3$ . The results quickly follow.  $\square$

We can also produce examples in  $G_2$  manifolds with an isometric involution.

**Proposition 6.6.** *Let  $(M, \varphi)$  be a  $G_2$  manifold with an isometric involution  $\sigma \neq \operatorname{id}$  such that  $\sigma^* \varphi = \varphi$  or  $\sigma^* \varphi = -\varphi$ . Then  $\operatorname{Fix}(\sigma)$  is an associative or coassociative submanifold in  $M$  respectively, if it is non-empty.*

We also have explicit examples of associatives and coassociatives.

**Example.** The first explicit examples of associatives in  $\mathbb{R}^7$  not arising from other geometries are given in [52] from symmetry and evolution equation considerations.

The first explicit non-trivial examples of coassociatives in  $\mathbb{R}^7$  are given in [20]. There are two dilation families: one which has one end asymptotic to a cone  $C$  on a non-round  $\mathcal{S}^3$ , and one which has two ends asymptotic to  $C \cup \mathbb{R}^4$ . The cone  $C$  was discovered earlier by Lawson–Osserman [50] and was the first example of a volume-minimizing submanifold which is not smooth (it is Lipschitz but not  $C^1$ ).

**Example.** In the Bryant–Salamon holonomy  $G_2$  metric on the spinor bundle of  $\mathcal{S}^3$  [5], the base  $\mathcal{S}^3$  is associative. In the Bryant–Salamon holonomy  $G_2$  metrics on  $\Lambda_+^2 T^* \mathcal{S}^4$  and  $\Lambda_+^2 T^* \mathbb{C}\mathbb{P}^2$  [5], the bases  $\mathcal{S}^4$  and  $\mathbb{C}\mathbb{P}^2$  are coassociative.

We now want to understand deformations of associatives and coassociatives, from which perturbation or gluing results will follow. We begin with associatives.

Notice that if  $P$  is an associative plane,  $u \in P$  and  $v \in P^\perp$  then  $u \times v \in P^\perp$  since  $\varphi(w, u, v) = g(w, u \times v) = g(v, w \times u) = 0$  for all  $w \in P$  since  $w \times u \in P$ . Thus, if  $N$  is associative, cross product gives a (Clifford) multiplication  $m : C^\infty(T^*N \otimes \nu(N)) \rightarrow C^\infty(\nu(N))$  (viewing tangent vectors as cotangent vectors via the metric). Hence, using the normal connection  $\nabla^\perp : C^\infty(\nu(N)) \rightarrow C^\infty(T^*N \otimes \nu(N))$  on  $\nu(N)$  we get a linear operator

$$\not{D} = m \circ \nabla^\perp : C^\infty(\nu(N)) \rightarrow C^\infty(\nu(N)).$$

We call  $\not{D}$  the *Dirac operator*. We see that its principal symbol is given by  $\sigma_{\not{D}}(x, \xi)v = i\xi \times v$ , so  $\not{D}$  is elliptic, and we also have that  $\not{D}^* = \not{D}$ .

**Remark.** Since a 3-manifold is always spin, we have a spinor bundle  $\mathbb{S}$  on  $N$ , a connection  $\nabla : C^\infty(\mathbb{S}) \rightarrow C^\infty(T^*M \otimes \mathbb{S})$  (a lift of the Levi-Civita connection) and we have Clifford multiplication  $m : C^\infty(T^*M \otimes \mathbb{S}) \rightarrow C^\infty(\mathbb{S})$  given by  $m(\xi, v) = \xi \cdot v$ . Hence we have a composition  $\not{D} = m \circ \nabla : C^\infty(\mathbb{S}) \rightarrow C^\infty(\mathbb{S})$ , which is a first order linear differential operator called the Dirac operator. Locally it is given by  $\not{D}v = \sum_i e_i \cdot \nabla_{e_i} v$ , so we have that  $\sigma_{\not{D}}(\xi, v) = i\xi \cdot v$ . Hence  $\not{D}$  is elliptic. Moreover  $\not{D}$  is self-adjoint.

In fact, it is possible (see e.g. [66]) to see that the complexified normal bundle  $\nu(N) \otimes \mathbb{C} = \mathbb{S} \otimes V$  for a  $\mathbb{C}^2$ -bundle  $V$  over  $N$ , so that the Dirac operator on  $\nu(N)$  is just a “twist” of the usual Dirac operator on  $\mathbb{S}$ .

Consider a compact associative  $N$ . We want to describe the associative deformations of  $N$ , just as in the case of special Lagrangians above. To be consistent with that previous setting, we will now use  $P$  to denote a nonlinear deformation map: we trust that this will not cause confusion given the context.

We know that  $\exp_v(N) = N_v$ , which is the graph of  $v$ , is associative for a normal vector field  $v$  if and only if  $*\exp_v^*(\chi) \in C^\infty(TM|_N)$  is 0. In fact, it turns out that  $P(v) = *\exp_v^*(\chi) \in C^\infty(\nu(N))$  since  $N$  is associative and

$$L_0P(v) = *d(v \lrcorner \chi) = \not{D}v.$$

Here  $L_0P$  is not typically surjective so we cannot apply our Implicit Function Theorem, except when  $\text{Ker } \not{D} = \text{Ker } \not{D}^* = \{0\}$ . However, we can still say something in these circumstances, for which we make a small digression to a more general situation.

Suppose  $X, Y$  are Banach spaces. Let  $U \subseteq X$  be an open set with  $0 \in U$  and let  $P : U \rightarrow Y$  be a smooth map with  $P(0) = 0$  such that  $L_0P : X \rightarrow Y$  is Fredholm.

Let  $\mathcal{I} = \text{Ker } L_0P$  and let  $\mathcal{O}$  be such that  $Y = L_0P(X) \oplus \mathcal{O}$ , which exists and is finite-dimensional by the assumption that  $L_0P$  is Fredholm. We then let  $Z = X \oplus \mathcal{O}$  and define  $F : U \oplus \mathcal{O} \rightarrow Y$  by  $F(u, y) = P(u) + y$ . We see that  $L_0F : X \oplus \mathcal{O} \rightarrow Y = L_0P(X) \oplus \mathcal{O}$  is given by  $L_0F(x, y) = L_0P(x) + y$  which is surjective and  $L_0F(x, y) = 0$  if and only if  $L_0P(x) = 0$  and  $y = 0$ , thus  $\text{Ker } L_0F = \text{Ker } L_0P \times \{0\}$ .

There exists  $W \subseteq X$  such that  $\text{Ker } L_0P \oplus W = X$ . Applying the Implicit Function Theorem, there exist open sets  $U_1 \subseteq \text{Ker } L_0P$  containing 0,  $U_2 \subseteq W$  containing 0 and  $U_3 \subseteq \mathcal{O}$  containing 0 and smooth maps  $G_2 : U_1 \rightarrow U_2$ ,  $G_3 : U_1 \rightarrow U_3$  such that

$$F^{-1}(0) \cap U_1 \times U_2 \times U_3 = \{(u, G_2(u), G_3(u)) : u \in U_1\}.$$

We also know that  $P(x) = 0$  if and only if  $F(x, y) = 0$  and  $y = 0$ . Hence

$$P^{-1}(0) \cap U_1 \times U_2 = \{(u, G_2(u)) : u \in G_3^{-1}(0)\}.$$

Let  $\mathcal{U} = U_1$  and define  $\pi : \mathcal{U} \rightarrow \mathcal{O}$  by  $\pi(u) = G_3(u)$ . Then  $P^{-1}(0) \cap U_1 \times U_2$  is a graph over  $\pi^{-1}(0)$ , and hence  $P^{-1}(0)$  is locally homeomorphic to  $\pi^{-1}(0)$ .

Sard's Theorem says that generically  $\pi^{-1}(y)$  is a smooth manifold of dimension  $\dim \mathcal{I} - \dim \mathcal{O} = \dim \text{Ker } L_0P - \dim \text{Coker } L_0P$ , which is the index of  $L_0P$ . Hence, the expected dimension of  $P^{-1}(0)$  is the index of  $L_0P$ .

In the associative setting we have that the linearisation is  $\not{D}$ , which is elliptic and thus Fredholm, and we know that  $\text{index } \not{D} = \dim \text{Ker } \not{D} - \dim \text{Ker } \not{D}^* = 0$ . We deduce the following [66].

**Theorem 6.7.** *The expected dimension of the moduli space of deformations of a compact associative 3-fold  $N$  in a  $G_2$  manifold is 0 and infinitesimal deformations of  $N$  are given by the kernel of  $\not{D}$  on  $\nu(N)$ . Moreover, if  $\text{Ker } \not{D} = \{0\}$  then  $N$  is rigid.*

**Remark.** The dimension of the kernel of  $\not{D}$  typically depends on the metric on  $N$  rather than just the topology, so it is usually difficult to determine. However, there are some cases where one can ensure the moduli space is smooth cf. [15].

**Example.** For the associative  $N = \mathcal{S}^3$  in  $\mathbb{S}(\mathcal{S}^3)$ ,  $\nu(N) = \mathbb{S}(\mathcal{S}^3)$  so  $\not{D}$  is just the usual Dirac operator. A theorem of Lichnerowicz states that  $\text{Ker } \not{D} = \{0\}$  as  $\mathcal{S}^3$  has positive scalar curvature so  $N$  is rigid.

**Example.** Corti–Haskins–Nordström–Pacini construct rigid associative  $\mathcal{S}^1 \times \mathcal{S}^2$ s in compact holonomy  $G_2$  twisted connected sums [10].

For coassociatives, the deformation theory is much better behaved, like for special Lagrangians [66].

**Theorem 6.8.** *Let  $N$  be a compact coassociative in a  $G_2$  manifold (or just a 7-manifold with closed  $G_2$  structure). The moduli space of deformations of  $N$  is a smooth manifold of dimension  $b_+^2(N)$ .*

*Proof.* Since  $N$  is coassociative the map  $v \mapsto v \lrcorner \varphi = \alpha_v$  defines an isomorphism from  $\nu(N)$  to a rank 3 vector bundle on  $N$ , which is  $\Lambda_+^2 T^*N$ , the 2-forms on  $N$  which are self-dual (so  $*\alpha = \alpha$ ). We can therefore view nearby submanifolds to  $N$  as graphs of self-dual 2-forms.



We know that  $N_v = \exp_v(N)$  is coassociative if and only if  $\exp_v^*(\varphi) = 0$ . We see that

$$\frac{d}{dt} \exp_{tv}^*(\varphi)|_{t=0} = \mathcal{L}_v \varphi = d(v \lrcorner \varphi) = d\alpha_v.$$

Hence nearby coassociatives  $N'$  to  $N$  are given by the zeros of  $P(\alpha) = d\alpha + Q(\alpha, \nabla\alpha)$ . Moreover, since  $\varphi = 0$  on  $N$ ,  $[\varphi] = 0$  on  $N'$  and hence  $P : C^\infty(\Lambda_+^2 T^*N) \rightarrow d(C^\infty(\Lambda^2 T^*N))$ .

Here  $P$  is not elliptic, but  $L_0 P = d$  has finite-dimensional kernel, the closed self-dual 2-forms, since  $d\alpha = 0$  implies that  $d^*\alpha = - * d * \alpha = 0$  so  $\alpha$  is harmonic. Moreover,  $L_0 P$  has injective symbol so it is overdetermined elliptic, which means that elliptic regularity still holds. Another way to deal with this is to consider  $F(\alpha, \beta) = P(\alpha) + d^*\beta$  for  $\beta$  a 4-form. Now  $F^{-1}(0)$  is the disjoint union of  $P^{-1}(0)$  and multiples of the volume form, as exact and coexact forms are orthogonal. Moreover,  $L_0 F(\alpha, \beta) = d\alpha + d^*\beta$  is now elliptic. Overall, we can apply our standard Implicit Function Theorem if we know that

$$d(C^{k+1,a}(\Lambda_+^2 T^*N)) = d(C^{k+1,a}(\Lambda^2 T^*N)).$$

This is true because by Hodge theory if  $\alpha$  is a 2-form, we can write  $\alpha = d^*\beta + \gamma$  for a 3-form  $\beta$  and a closed form  $\gamma$ , so  $d\alpha = dd^*\beta = d(d^*\beta + *d^*\beta)$  and  $d^*\beta + *d^*\beta$  is self-dual.  $\square$

**Example.** The  $S^4$  and  $\mathbb{C}\mathbb{P}^2$  in the Bryant–Salamon metrics on  $\Lambda_+^2 T^*S^4$  and  $\Lambda_+^2 T^*\mathbb{C}\mathbb{P}^2$  have  $b_+^2 = 0$  and so are rigid.

For a K3 surface and  $T^4$  we have  $b_+^2 = 3$  and  $\Lambda_+^2$  is trivial, so we can hope to find coassociative K3 and  $T^4$  fibrations of compact  $G_2$  manifolds. There is a programme [47] for constructing a coassociative K3 fibration (with singular fibres). Towards completing this programme, the first examples of compact coassociative 4-folds with conical singularities in compact holonomy  $G_2$  twisted connected sums were constructed in [61].

Again, we have a similar isometric embedding result for coassociative 4-folds, motivated by the deformation theory result [3].

**Theorem 6.9.** *Any compact oriented real analytic Riemannian 4-manifold whose bundle of self-dual 2-forms is trivial can be isometrically embedded in a  $G_2$  manifold as the fixed points of an isometric involution.*

**Remark.** The deformation theory results for compact associative and coassociative submanifolds have been extended to certain non-compact, singular and boundary settings, for example in [16, 44, 48, 55, 56, 58].

## 7 Cayley submanifolds

We now discuss our final class of calibrated submanifolds.

**Theorem 7.1.** *On a  $\text{Spin}(7)$  manifold  $(M^8, \Phi)$  (so  $\Phi$  is a closed admissible form),  $\Phi$  is a calibration.*

*Proof.* Let  $P$  be a plane in  $\mathbb{R}^8 \cong \mathbb{C}^4$ . Since  $\text{SU}(4) \subseteq \text{Spin}(7)$ , by the proof of Wirtinger's inequality (Theorem 3.2), we can choose  $A \in \text{Spin}(7)$  so that  $A(P)$  is spanned by

$$\{e_1, \cos\theta_1 e_2 + \sin\theta_1 e_3, e_5, \cos\theta_2 e_6 + \sin\theta_2 e_7\}.$$

We take the standard  $\text{Spin}(7)$  form  $\Phi$  on  $\mathbb{R}^8$  to be:

$$\begin{aligned} \Phi = & dx_{1234} + dx_{1256} + dx_{1278} + dx_{1357} - dx_{1367} - dx_{1458} - dx_{1467} \\ & + dx_{5678} + dx_{3478} + dx_{3456} + dx_{2468} - dx_{2457} - dx_{2367} - dx_{2358}, \end{aligned}$$

again using the notation  $dx_{ij\dots k} = dx_i \wedge dx_j \wedge \dots \wedge dx_k$ . Then  $\Phi|_P = (\cos\theta_1 \cos\theta_2 + \sin\theta_1 \sin\theta_2) \text{vol}_P = \cos(\theta_1 - \theta_2) \text{vol}_P$ . Hence  $\Phi$  is a calibration as it is closed.  $\square$

We can thus define our calibrated submanifolds in  $\text{Spin}(7)$  manifolds.

**Definition 7.2.** Submanifolds in  $(M^8, \Phi)$  calibrated by  $\Phi$  are called Cayley 4-folds.

**Remark.** The name Cayley submanifolds is because of the relation between the submanifolds and the octonions or Cayley numbers  $\mathbb{O}$ .

We can relate Cayley submanifolds to all of the other calibrated geometries we have seen.

**Proposition 7.3.** (a) *Complex surfaces and special Lagrangian 4-folds in  $\mathbb{C}^4$  are Cayley in  $\mathbb{R}^8 = \mathbb{C}^4$ .*

(b) *Write  $\mathbb{R}^8 = \mathbb{R} \times \mathbb{R}^7$ . Then  $\mathbb{R} \times S$  is Cayley if and only if  $S$  is associative in  $\mathbb{R}^7$  and  $N \subseteq \mathbb{R}^7$  is Cayley in  $\mathbb{R}^8$  if and only if  $N$  is coassociative in  $\mathbb{R}^7$ .*

*Proof.* Recall the Kähler form  $\omega$  and holomorphic volume form  $\Upsilon$  on  $\mathbb{C}^4$  and the  $G_2$  3-form  $\varphi$  on  $\mathbb{R}^7$ .

Part (a) is immediate from the formula  $\Phi = \frac{1}{2}\omega^2 + \text{Re } \Upsilon$ , since complex surfaces are calibrated by  $\frac{1}{2}\omega^2$ , special Lagrangians are calibrated by  $\text{Re } \Upsilon$ ,  $\Upsilon$  vanishes on complex surfaces and  $\omega$  vanishes on special Lagrangians.

Part (b) follows immediately from the formula  $\Phi = dx_1 \wedge \varphi + *\varphi$ .  $\square$

We can also use an isometric involution to construct Cayley submanifolds as in our previous calibrated geometries.

**Proposition 7.4.** *Let  $(M, \Phi)$  be a  $\text{Spin}(7)$  manifold and let  $\sigma \neq \text{id}$  be an isometric involution with  $\sigma^*\Phi = \Phi$ . Then  $\text{Fix}(\sigma)$  is Cayley submanifold, if it is non-empty.*

**Example.** The first interesting explicit examples of Cayleys in  $\mathbb{R}^8$  not arising from other geometries were given in [53] and are asymptotic to cones.

**Example.** The base  $\mathcal{S}^4$  in the Bryant–Salamon holonomy  $\text{Spin}(7)$  metric on  $\mathbb{S}_+(\mathcal{S}^4)$  [5] is Cayley.

We now discuss deformations of a compact Cayley  $N$ , for which we need some discussion of algebra related to  $\text{Spin}(7)$ . Since  $\Lambda^2(\mathbb{R}^8)^*$  is 28-dimensional and the 21-dimensional Lie algebra of  $\text{Spin}(7)$  sits inside the space of 2-forms, we must have a distinguished 7-dimensional subspace  $\Lambda_7^2$  of 2-forms on  $\mathbb{R}^8$ . So what is this subspace? Let  $u, v \in \mathbb{R}^8$ . Then we can construct a 2-form  $u \wedge v$ , viewing  $u, v$  as cotangent vectors. We can also construct a 2-form from  $u, v$  by considering  $\Phi(u, v, \cdot, \cdot)$ . It is then true that

$$\Lambda_7^2 = \{u \wedge v + \Phi(u, v, \cdot, \cdot) : u, v \in \mathbb{R}^8\}.$$

When  $P$  is a Cayley plane and  $u, v \in P$  are orthogonal we see that  $\Phi(u, v, \cdot, \cdot) = *_{P}(u \wedge v)$  so that  $u \wedge v + \Phi(u, v, \cdot, \cdot)$  is self-dual on  $P$ . Since  $\Lambda_+^2 P^*$  is 3-dimensional, we see that there must be a 4-dimensional space  $E$  of 2-forms on  $P$  such that  $\Lambda_7^2|_P = \Lambda_+^2 P^* \oplus E$ . Moreover, if  $u \in P$  and  $v \in P^\perp$  then  $m(u, v) = u \wedge v + \Phi(u, v, \cdot, \cdot) \in E$  and the map  $m : P \times P^\perp \rightarrow E$  is surjective.

Now let us move to a Cayley submanifold  $N$  in a  $\text{Spin}(7)$  manifold  $(M, \Phi)$ . On  $M$  we have a rank 7 bundle  $\Lambda_7^2$  of 2-forms and we have that  $\Lambda_7^2|_N = \Lambda_+^2 T^*N \oplus E$  for some rank 4 bundle  $E$  over  $N$ . The map  $m$  above defines a (Clifford) multiplication  $m : C^\infty(T^*N \otimes \nu(N)) \rightarrow C^\infty(E)$  (viewing tangent vectors as cotangent vectors via the metric), and thus using the normal connection  $\nabla^\perp : C^\infty(\nu(N)) \rightarrow C^\infty(T^*N \otimes \nu(N))$  we get a linear first order differential operator

$$\mathcal{D}_+ = m \circ \nabla^\perp : C^\infty(\nu(N)) \rightarrow C^\infty(E).$$

Again this an elliptic operator called the *positive Dirac operator*, but it is not self-adjoint: its adjoint is the negative Dirac operator from  $E$  to  $\nu(N)$ .

**Remark.** If  $N$  is spin, the spinor bundle  $\mathbb{S}$  splits as  $\mathbb{S}_+ \oplus \mathbb{S}_-$ , and the Dirac operator  $\mathcal{D}$  splits into  $\mathcal{D}_\pm$  from  $\mathbb{S}_\pm$  to  $\mathbb{S}_\mp$  so that  $\mathcal{D}(v_+, v_-) = (\mathcal{D}_- v_-, \mathcal{D}_+ v_+)$ . Hence  $\mathcal{D}^* = \mathcal{D}$  says that  $\mathcal{D}_\pm^* = \mathcal{D}_\mp$ .

It turns out (see, for example, [66]) that there exists a  $\mathbb{C}^2$ -bundle  $V$  on  $N$  so that  $\nu(N) \otimes \mathbb{C} = \mathbb{S}_+ \otimes V$ ,  $E \otimes \mathbb{C} = \mathbb{S}_- \otimes V$  and  $\mathcal{D}_+$  on  $\nu(N)$  is a “twist” of the usual positive Dirac operator. However, not every 4-manifold is spin, so we cannot always make this identification.

On  $\mathbb{O}$  there exists a 4-fold cross product, whose real part gives  $\Phi$  and whose imaginary part we call  $\tau$ . Perhaps unsurprisingly, we have the following result, which we will leave as an exercise for the reader.

**Lemma 7.5.** *A 4-plane  $P$  in  $\mathbb{R}^8$  satisfies  $\tau|_P \equiv 0$  if and only if it admits an orientation so that it is calibrated by  $\Phi$ .*

We can extend  $\tau$  to a  $\text{Spin}(7)$  manifold, except that we need a rank 7 vector bundle on  $M$  in which  $\tau$  takes values: we have one, namely  $\Lambda_7^2$ . So we have the following alternative characterisation of Cayley 4-folds.

**Lemma 7.6.** *A submanifold  $N$  in a  $\text{Spin}(7)$  manifold is Cayley (up to a choice of orientation) if and only if  $\tau \in C^\infty(\Lambda^4 T^*M; \Lambda_7^2)$  vanishes on  $N$ .*

Now suppose that  $N$  is a compact Cayley 4-fold. Then the zeros of the equation  $F(v) = * \exp_v^*(\tau)$  for  $v \in C^\infty(\nu(N))$  define Cayley deformations (as the graph of  $v$ ). We know that  $F$  takes values in  $\Lambda_7^2|_N = \Lambda_+^2 T^*N \oplus E$  and it turns out that

$$L_0 F(v) = *d(v \lrcorner \tau) = \not{D}_+ v$$

since  $N$  is Cayley. So, we potentially have a problem because  $F$  does not necessarily take values only in  $E$  (and in general it will not just take values in  $E$ ). However, the Cayley condition on  $N$  means that  $F(v) = 0$  if and only if  $P(v) = \pi_E F(v) = 0$ , where  $\pi_E$  is the projection onto  $E$  (again, we are using  $P$  to denote the nonlinear deformation map as in our previous discussion, and we expect it will not cause confusion given the context). Then the operator  $P : C^\infty(\nu(N)) \rightarrow C^\infty(E)$  and  $L_0 P = \not{D}_+$  is elliptic.

Again, we cannot say that  $L_0 P$  is surjective, so we have the following using the same argument as in the lead up to Theorem 6.7, cf. [66].

**Theorem 7.7.** *The expected dimension of the moduli space of deformations of a compact Cayley 4-fold  $N$  in a  $\text{Spin}(7)$  manifold is  $\text{ind } \not{D}_+ = \dim \text{Ker } \not{D}_+ - \dim \text{Ker } \not{D}_+^*$  with infinitesimal deformations given by  $\text{Ker } \not{D}_+$  on  $\nu(N)$ . Moreover,*

$$\text{ind } \not{D}_+ = \frac{1}{2} \sigma(N) + \frac{1}{2} \chi(N) - [N].[N],$$

where  $\sigma(N) = b_+^2(N) - b_-^2(N)$  (the signature of  $N$ ),  $\chi(N) = 2b^0(N) - 2b^1(N) + b^2(N)$  (the Euler characteristic of  $N$ ) and  $[N].[N]$  is the self-intersection of  $N$ , which is the Euler number of  $\nu(N)$ .

**Example.** For the Cayley  $N = \mathcal{S}^4$  in  $\mathbb{S}_+(\mathcal{S}^4)$ ,  $\nu(N) = \mathbb{S}_+(\mathcal{S}^4)$  and  $\not{D}_+$  is the usual positive Dirac operator. Again, since  $N$  has positive scalar curvature, we see that  $\text{Ker } \not{D}_\pm = \{0\}$  so  $N$  is rigid.

**Remark.** Theorem 7.7 has been extended to various other non-compact, singular and boundary settings, for example in [68, 70, 71].

## 8 The angle theorem

To conclude these notes, we now discuss a very natural and elementary problem in Euclidean geometry where calibrations play a major, and perhaps unexpected, role.

If one takes two lines in  $\mathbb{R}^2$  intersecting transversely, then their union is never length-minimizing. A natural question to ask is: does this persist in higher dimensions? In other words, when is the union of two transversely intersecting  $n$ -planes in  $\mathbb{R}^{2n}$  volume-minimizing? Two such planes are determined by the  $n$  angles between them as follows.

**Lemma 8.1.** *Let  $P, Q$  be oriented  $n$ -planes in  $\mathbb{R}^{2n}$ . There exists an orthonormal basis  $e_1, \dots, e_{2n}$  for  $\mathbb{R}^{2n}$  such that  $P = \text{Span}\{e_1, \dots, e_n\}$  and*

$$Q = \text{Span}\{\cos \theta_1 e_1 + \sin \theta_1 e_{n+1}, \dots, \cos \theta_n e_n + \sin \theta_n e_{2n}\}$$

where  $0 \leq \theta_1 \leq \dots \leq \theta_{n-1} \leq \frac{\pi}{2}$  and  $\theta_{n-1} \leq \theta_n \leq \pi - \theta_{n-1}$ . These angles are called the characterising angles of  $P, Q$ .

*Proof.* The proof is very similar to the argument in the proof of Wirtinger's inequality (Theorem 3.2). Choose unit  $e_1 \in P$  and maximise  $\langle e_1, u_1 \rangle$  for  $u_1 \in Q$ , and let  $e_{n+1} \in P^\perp$  be defined by  $u_1 = \cos \theta_1 e_1 + \sin \theta_1 e_{n+1}$ . Now choose  $e_2 \in P \cap e_1^\perp$  and maximise  $\langle e_2, u_2 \rangle$  for  $u_2 \in Q \cap u_1^\perp$ , then proceed by induction.  $\square$

If the characterising angles of  $P, Q$  are  $\theta_1, \dots, \theta_n$ , then the characterising angles of  $P, -Q$  are  $\psi_1, \dots, \psi_n$  where  $\psi_j = \theta_j$  for  $j = 1, \dots, n-1$  and  $\psi_n = \pi - \theta_n$ .

The idea of the following theorem is that the union of  $P \cup Q$  is area-minimizing if  $P, -Q$  are not too close together [49].

**Theorem 8.2** (Angle Theorem). *Let  $P, Q$  be oriented transverse  $n$ -planes in  $\mathbb{R}^{2n}$  and let  $\psi_1, \dots, \psi_n$  be the characterising angles between  $P, -Q$ . Then  $P \cup Q$  is volume-minimizing if and only if  $\psi_1 + \dots + \psi_n \geq \pi$ .*

Notice that this criteria is impossible to fulfill in 1 dimension.

*Proof.* We will sketch the proof which involves calibrations in a fundamental way in both directions. For details, look at [19].

First if  $P \cup Q$  does not satisfy the angle condition, we can choose coordinates by Lemma 8.1 so that  $P = P(-\frac{\psi}{2})$  and  $-Q = P(\frac{\psi}{2})$  where  $\psi = (\psi_1, \dots, \psi_n)$  and  $P(\psi) = \{(e^{i\psi_1}x_1, \dots, e^{i\psi_n}x_n) : (x_1, \dots, x_n) \in \mathbb{R}^n\}$  as given earlier. We know that we have a special Lagrangian Lawlor neck  $N$  asymptotic to  $P(-\frac{\psi'}{2}) \cup P(\frac{\psi'}{2})$  for any  $\psi'$  where  $\sum_{i=1}^n \psi'_i = \pi$ . The claim is then that since  $\sum \psi_i < \pi$  we can find  $\psi'$  so that  $\sum \psi'_i = \pi$  and  $N \cap P(\pm \frac{\psi'}{2})$  is non-empty (in fact, an ellipsoid). This is actually a way to characterise  $N$ . Hence since  $N$  is calibrated by  $\text{Im } \Upsilon$  and  $\text{Im } \Upsilon|_{P \cup Q} < \text{vol}_{P \cup Q}$ ,  $P \cup Q$  cannot be volume-minimizing by the usual Stokes' Theorem argument for calibrated submanifolds.

We now provide a few extra details, for which we need to describe  $N$ . For maps  $z_1, \dots, z_n : \mathbb{R} \rightarrow \mathbb{C}$  define

$$N = \{(t_1 z_1(s), \dots, t_n z_n(s)) \in \mathbb{C}^n : s \in \mathbb{R}, t_1, \dots, t_n \in \mathbb{R}, \sum_{j=1}^n t_j^2 = 1\}.$$

It is not difficult to calculate that  $N$  is special Lagrangian with phase  $i$  (so calibrated by  $\text{Im } \Upsilon$ ) if and only if

$$\frac{dz_j}{z_j} \frac{ds}{ds} = i f_j \overline{z_1 \dots z_n}$$

for positive real functions  $f_j$ .

Suppose that  $f_j = 1$  for all  $j$ . Write  $z_j = r_j e^{i\theta_j}$ , let  $\theta = \sum_{j=1}^n \theta_j$  and suppose that  $z_j(0) = c_j > 0$ . From the differential equation, one quickly sees that  $r_j^2 = c_j^2 + u$  for some function  $u$  with  $u(0) = 0$  and  $r_1 \dots r_n \cos \theta = c_1 \dots c_n$ .

If we now suppose that  $u = t^2$ , we see that

$$\theta_j(t) = \int_0^t \frac{a_j dt}{(1 + a_j t^2) \sqrt{\frac{1}{t^2} ((1 + a_1 t^2) \dots (1 + a_n t^2) - 1)}}$$

where  $a_j = c_j^{-2}$ . We observe that  $\theta \rightarrow \pm \frac{\pi}{2}$  as  $t \rightarrow \pm \infty$  and hence  $N$ , which is a Lawlor neck, is asymptotic to a pair of planes where the sum of the angles is  $\pm \frac{\pi}{2}$ .

Now fix  $t > 0$  and define

$$f : X = \{(a_1, \dots, a_n) \in \mathbb{R}^n : a_j \geq 0\} \rightarrow Y = \{(\theta_1, \dots, \theta_n) \in \mathbb{R}^n : \theta_j \geq 0, \sum_{j=1}^n \theta_j < \frac{\pi}{2}\}$$

by  $f(a_1, \dots, a_n) = (\theta_1, \dots, \theta_n)$  where

$$\theta_j = \int_0^t \frac{a_j dt}{(1 + a_j t^2) \sqrt{\frac{1}{t^2} ((1 + a_1 t^2) \dots (1 + a_n t^2) - 1)}}.$$

It is clear that if  $n = 1$ ,  $f : X \rightarrow Y$  is surjective. We want to show it is surjective for all  $n$ . For  $\theta \in (0, \frac{\pi}{2})$  define  $H_\theta = \{(\theta_1, \dots, \theta_n) \in Y : \sum_{j=1}^n \theta_j = \theta\}$ . By our discussion above we see that

$$f^{-1}(H_\theta) \subseteq S_\theta = \{(a_1, \dots, a_n) \in X : (1 + a_1 t^2) \dots (1 + a_n t^2) = \cos^{-2} t\}.$$

Notice that if the degree of  $f : \partial S_\theta \rightarrow \partial H_\theta$  is 1 then the degree of  $f : S_\theta \rightarrow H_\theta$  is 1. Thus, by induction on  $n$ , we see that  $f : S_\theta \rightarrow H_\theta$  is surjective.

Now, given any plane  $\{(e^{i\theta_1}x_1, \dots, e^{i\theta_n}x_n) : (x_1, \dots, x_n) \in \mathbb{R}^n\}$  where  $(\theta_1, \dots, \theta_n) \in Y$ ,  $\theta_j \neq 0$  for all  $j$ , we see that we can choose a Lawlor neck  $N$  which intersects the plane in a hypersurface as claimed.

We now move to the other direction in the statement of the Angle Theorem. If  $P \cup Q$  does satisfy the angle condition, then (by choosing coordinates so that  $P = \mathbb{R}^n$  and  $Q$  is in standard position) we claim that it is calibrated by a so-called *Nance calibration*:

$$\eta(u_1, \dots, u_n) = \operatorname{Re}((dx_1 + u_1 dy_1) \wedge \dots \wedge (dx_n + u_n dy_n))$$

where  $u_1, \dots, u_n \in \mathcal{S}^2 \subseteq \operatorname{Im}\mathbb{H}$ . If  $u_m = i$  for all  $m$  then  $\eta = \operatorname{Re} \Upsilon$ , so it is believable that it is a calibration in general, but we now show that it is indeed true.

Let  $x_1, y_1, \dots, x_n, y_n$  be coordinates on  $\mathbb{R}^{2n}$ . We call an  $n$ -form  $\eta$  on  $\mathbb{R}^{2n}$  a torus form if  $\eta$  lies in the span of forms of type

$$dx_{i_1} \wedge \dots \wedge dx_{i_k} \wedge dy_{j_1} \wedge \dots \wedge dy_{j_l}$$

where  $\{i_1, \dots, i_k\} \cap \{j_1, \dots, j_l\} = \emptyset$  and  $\{i_1, \dots, i_k\} \cup \{j_1, \dots, j_l\} = \{1, \dots, n\}$ . We now claim that a torus form  $\eta$  is a calibration if and only if

$$\eta(\cos \theta_1 e_1 + \sin \theta_1 e_{n+1}, \dots, \cos \theta_n e_n + \sin \theta_n e_{2n}) \leq 1$$

for all  $\theta_1, \dots, \theta_n \in \mathbb{R}$ .

For  $n = 1$ ,  $\eta = dx_1 \wedge dy_1$  which is a calibration. Suppose that the result holds for  $n = k$ . Let  $\eta$  be a torus form on  $\mathbb{R}^{2(k+1)}$  and rescale  $\eta$  so that the maximum of  $\eta$  is 1 and is attained at some plane. The idea is to show using the argument in the proof of Wirtinger's inequality to put planes in standard position that we can write  $\eta = e_1 \wedge \eta_1 + e_2 \wedge \eta_2$  where  $e_1, e_2$  are orthonormal and span an  $\mathbb{R}^2$  and  $\eta_1, \eta_2$  are torus forms on  $\mathbb{R}^{2k}$ . The claim then follows by induction on  $n$ .

Hence, the Nance calibration  $\eta$  above is a calibration and moreover we know  $P(\theta)$  is calibrated by  $\eta(u)$  if and only if

$$\prod_{j=1}^n (\cos \theta_j + \sin \theta_j u_j) = 1.$$

We then just need to find the  $u_j$  determined by  $\theta_j$ . Notice that the condition that  $\psi_1 + \dots + \psi_n \geq \pi$  holds if and only if  $\theta_n \leq \theta_1 + \dots + \theta_{n-1}$ . If we write  $\cos \theta_j + \sin \theta_j u_j = w_j \bar{w}_{j+1}$  where  $w_{n+1} = w_1$  and  $w_j$  are unit imaginary quaternions then the product condition is satisfied and we just need  $\langle w_j, w_{j+1} \rangle = \cos \theta_j$ , which is equivalent to finding  $n$  points on the unit 2-sphere so that  $d(w_j, w_{j+1}) = \theta_j$ , where  $\theta_n \leq \theta_1 + \dots + \theta_{n-1}$ . This is indeed possible, by considering an  $n$ -sided spherical polygon.  $\square$

## References

- [1] S. Brendle, *Embedded minimal tori in  $S^3$  and the Lawson conjecture*, Acta Math. **211** (2013), no. 2, 177–190.
- [2] R. L. Bryant, *Submanifolds and special structures on the octonians*, J. Differential Geom. **17** (1982), 185–232.
- [3] R. L. Bryant, *Calibrated embeddings in the special Lagrangian and coassociative cases*, Ann. Global Anal. Geom. **18** (2000), no. 3-4, 405–435.
- [4] R. L. Bryant, *Second order families of special Lagrangian 3-folds*, Perspectives in Riemannian geometry, pp. 63–98, CRM Proc. Lecture Notes **40**, Amer. Math. Soc., Providence, RI, 2006.
- [5] R. L. Bryant and S. Salamon, *On the construction of some complete metrics with exceptional holonomy*, Duke Math. J. **58** (1989), 829–850.
- [6] A. Butscher, *Deformations of minimal Lagrangian submanifolds with boundary*, Proc. Amer. Math. Soc. **131** (2002) 1953–1964.
- [7] A. Butscher, *Regularizing a singular special Lagrangian variety*, Comm. Anal. Geom. **12** (2004), 733–791.
- [8] E. Carberry, *Associative cones in the imaginary octonions*, Riemann surfaces, harmonic maps and visualization, pp. 249–263, OCAMI Stud. **3**, Osaka Munic. Univ. Press, Osaka, 2010.
- [9] E. Carberry and I. McIntosh, *Minimal Lagrangian 2-tori in  $\mathbb{C}\mathbb{P}^2$  come in real families of every dimension*, J. London Math. Soc. (2) **69** (2004), no. 2, 531–544.

- [10] A. Corti, M. Haskins, J. Nordström and T. Pacini, *G<sub>2</sub>-manifolds and associative submanifolds via semi-Fano 3-folds*, Duke Math. J. **164** (2015), no. 10, 1971–2092.
- [11] T. Eguchi and A. J. Hanson, *Asymptotically flat self-dual solutions to Euclidean gravity*, Phys. Lett. **74B** (3) (1978), 249–251.
- [12] D. Fox, *Second order families of coassociative 4-folds*, Thesis (Ph.D.) Duke University, ProQuest LLC, Ann Arbor, MI, 2005.
- [13] D. Fox, *Coassociative cones ruled by 2-planes*, Asian J. Math. **11** (2007), no. 4, 535–553.
- [14] D. Fox, *Cayley cones ruled by 2-planes: desingularization and implications of the twistor fibration*, Comm. Anal. Geom. **16** (2008), no. 5, 937–968.
- [15] D. Gayet, *Smooth moduli spaces of associative submanifolds*, Q. J. Math. **65** (2014), no. 4, 1213–1240.
- [16] D. Gayet and F. Witt, *Deformations of associative submanifolds with boundary*, Adv. Math. **226** (2011), no. 3, 2351–2370.
- [17] E. Goldstein, *Calibrated fibrations on noncompact manifolds via group actions*, Duke Math. J. **110** (2001), no. 2, 309–343.
- [18] A. Gray, E. Abbena and S. Salamon, *Modern differential geometry of curves and surfaces with Mathematica®*, Third edition, Studies in Advanced Mathematics, Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [19] R. Harvey, *Spinors and calibrations*, Perspectives in Mathematics **9**, Academic Press, Inc., Boston, MA, 1990.
- [20] R. Harvey and H. B. Lawson, *Calibrated geometries*, Acta Math. **148** (1982), 47–157.
- [21] M. Haskins, *Special Lagrangian cones*, Amer. J. Math. **126** (2004), no. 4, 845–871.
- [22] M. Haskins, *The geometric complexity of special Lagrangian T<sup>2</sup>-cones*, Invent. Math. **157** (2004), no. 1, 11–70.
- [23] M. Haskins and N. Kapouleas, *Special Lagrangian cones with higher genus links*, Invent. Math. **167** (2007), 223–294.
- [24] G. Huisken and T. Ilmanen, *The inverse mean curvature flow and the Riemannian Penrose inequality*, J. Differential Geom. **59** (2001), no. 3, 353–437.
- [25] Y. Imagi, D. D. Joyce and J. Oliveira dos Santos, *Uniqueness results for special Lagrangians and Lagrangian mean curvature flow expanders in  $\mathbb{C}^m$* , Duke Math. J. **165** (2016), no. 5, 847–933.
- [26] M. Ionel, *Second order families of special Lagrangian submanifolds in  $\mathbb{C}^4$* , J. Differential Geom. **65** (2003), no. 2, 211–272.
- [27] M. Ionel, S. Karigiannis and M. Min-Oo, *Bundle constructions of calibrated submanifolds in  $\mathbb{R}^7$  and  $\mathbb{R}^8$* , Math. Res. Lett. **12** (2005), no. 4, 493–512.
- [28] M. Ionel and M. Min-Oo, *Cohomogeneity one special Lagrangian 3-folds in the deformed and the resolved conifolds*, Illinois J. Math. **52** (2008), no. 3, 839–865.
- [29] K. Irie, F. Marques and A. Neves, *Density of minimal hypersurfaces for generic metrics*, Ann. of Math. (2) **187** (2018), no. 3, 963–972.
- [30] D. D. Joyce, *Evolution equations for special Lagrangian 3-folds in  $\mathbb{C}^3$* , Ann. Global Anal. Geom. **20** (2001), no. 4, 345–403.
- [31] D. D. Joyce, *Constructing special Lagrangian m-folds in  $\mathbb{C}^m$  by evolving quadrics*, Math. Ann. **320** (2001), no. 4, 757–797.
- [32] D. D. Joyce, *Ruled special Lagrangian 3-folds in  $\mathbb{C}^3$* , Proc. London Math. Soc. (3) **85** (2002), no. 1, 233–256.
- [33] D. D. Joyce, *Special Lagrangian m-folds in  $\mathbb{C}^m$  with symmetries*, Duke Math. J. **115** (2002), no. 1, 1–51.
- [34] D. D. Joyce, *U(1)-invariant special Lagrangian 3-folds in  $\mathbb{C}^3$  and special Lagrangian fibrations*, Turkish J. Math. **27** (2003), no. 1, 99–114.

- [35] D. D. Joyce, *Special Lagrangian submanifolds with isolated conical singularities. V. Survey and applications*, J. Differential Geom. **63** (2003), 279–347.
- [36] D. D. Joyce, *Special Lagrangian submanifolds with isolated conical singularities. II. Moduli spaces*, Ann. Global Anal. Geom. **25** (2004), 301–352.
- [37] D. D. Joyce, *Special Lagrangian submanifolds with isolated conical singularities. III. Desingularization, the unobstructed case*, Ann. Global Anal. Geom. **26** (2004), 1–58.
- [38] D. D. Joyce, *Special Lagrangian submanifolds with isolated conical singularities. IV. Desingularization, obstructions and families*, Ann. Global Anal. Geom. **26** (2004), 117–174.
- [39] D. D. Joyce,  *$U(1)$ -invariant special Lagrangian 3-folds. I. Nonsingular solutions*, Adv. Math. **192** (2005), no. 1, 35–71.
- [40] D. D. Joyce,  *$U(1)$ -invariant special Lagrangian 3-folds. II. Existence of singular solutions*, Adv. Math. **192** (2005), no. 1, 72–134.
- [41] D. D. Joyce,  *$U(1)$ -invariant special Lagrangian 3-folds. III. Properties of singular solutions*, Adv. Math. **192** (2005), no. 1, 135–182.
- [42] D. D. Joyce, *Riemannian holonomy groups and calibrated geometry*, Oxford Graduate Texts in Mathematics **12**, Oxford University Press, Oxford, 2007.
- [43] D. D. Joyce, *Special Lagrangian 3-folds and integrable systems*, Surveys on geometry and integrable systems, pp. 189–233, Adv. Stud. Pure Math. **51**, Math. Soc. Japan, Tokyo, 2008.
- [44] D. D. Joyce and S. Salur, *Deformations of asymptotically cylindrical coassociative submanifolds with fixed boundary*, Geom. Topol. **9** (2005), 1115–1146.
- [45] S. Karigiannis and N. C.-H. Leung, *Deformations of calibrated subbundles of Euclidean spaces via twisting by special sections*, Ann. Global Anal. Geom. **42** (2012), no. 3, 371–389.
- [46] S. Karigiannis and M. Min-Oo, *Calibrated subbundles in noncompact manifolds of special holonomy*, Ann. Global Anal. Geom. **28** (2005), no. 4, 371–394.
- [47] A. G. Kovalev, *Coassociative  $K3$  fibrations of compact  $G_2$ -manifolds*, arXiv:math/0511150.
- [48] A. Kovalev and J. D. Lotay, *Deformations of compact coassociative 4-folds with boundary*, J. Geom. Phys. **59** (2009), 63–73.
- [49] G. Lawlor, *The angle criterion*, Invent. Math. **95** (1989), 437–446.
- [50] H. B. Lawson and R. Osserman, *Non-existence, non-uniqueness and irregularity of solutions to the minimal surface system*, Acta Math. **139** (1977), 1–17.
- [51] Y.-I. Lee, *Embedded special Lagrangian submanifolds in Calabi–Yau manifolds*, Comm. Anal. Geom. **11** (2003), 391–423.
- [52] J. D. Lotay, *Constructing associative 3-folds by evolution equations*, Comm. Anal. Geom. **13** (2005), 999–1037.
- [53] J. D. Lotay, *2-ruled calibrated 4-folds in  $\mathbb{R}^7$  and  $\mathbb{R}^8$* , J. London Math. Soc. **74** (2006), 219–243.
- [54] J. D. Lotay, *Calibrated submanifolds of  $\mathbb{R}^7$  and  $\mathbb{R}^8$  with symmetries*, Q. J. Math. **58** (2007), 53–70.
- [55] J. D. Lotay, *Coassociative 4-folds with conical singularities*, Comm. Anal. Geom. **15** (2007), 891–946.
- [56] J. D. Lotay, *Deformation theory of asymptotically conical coassociative 4-folds*, Proc. London Math. Soc. **99** (2009), 386–424.
- [57] J. D. Lotay, *Desingularization of coassociative 4-folds with conical singularities*, Geom. Funct. Anal. **18** (2009), 2055–2100.
- [58] J. D. Lotay, *Asymptotically conical associative 3-folds*, Q. J. Math. **62** (2011), 131–156.
- [59] J. D. Lotay, *Ruled Lagrangian submanifolds of the 6-sphere*, Trans. Amer. Math. Soc. **363** (2011), 2305–2339.
- [60] J. D. Lotay, *Associative submanifolds of the 7-sphere*, Proc. London Math. Soc. **105** (2012), 1183–1214.
- [61] J. D. Lotay, *Stability of coassociative conical singularities*, Comm. Anal. Geom. **20** (2012), 803–867.

- 
- [62] J. D. Lotay, *Desingularization of coassociative 4-folds with conical singularities: obstructions and applications*, Trans. Amer. Math. Soc. **366** (2014), 6051–6092.
- [63] F. C. Marques and A. Neves, *Min-max theory and the Willmore conjecture*, Ann. of Math. (2) **179** (2014), no. 2, 683–782.
- [64] F. C. Marques and A. Neves, *Existence of infinitely many minimal hypersurfaces in positive Ricci curvature*, Invent. Math. **209** (2017), no. 2, 577–616.
- [65] I. McIntosh, *Special Lagrangian cones in  $\mathbb{C}^3$  and primitive harmonic maps*, J. London Math. Soc. (2) **67** (2003), no. 3, 769–789.
- [66] R. C. McLean, *Deformations of calibrated submanifolds*, Comm. Anal. Geom. **6** (1998), 705–747.
- [67] K. Moore, *Cayley deformations of compact complex surfaces*, arXiv:1710.08799.
- [68] K. Moore, *Deformations of conically singular Cayley submanifolds*, J. Geom. Anal. (2018).
- [69] C. B. Morrey, *Multiple Integrals in the Calculus of Variations*, Grundlehren Series Volume 130, Springer–Verlag, Berlin, 1966.
- [70] M. Ohst, *Deformations of compact Cayley submanifolds with boundary*, arXiv:1405.7886.
- [71] M. Ohst, *Deformations of asymptotically cylindrical Cayley submanifolds*, arXiv:1506.00110.
- [72] T. Pacini, *Special Lagrangian conifolds, I: moduli spaces*, Proc. London Math. Soc. (3) **107** (2013), 198–224.
- [73] T. Pacini, *Special Lagrangian conifolds, II: gluing constructions in  $\mathbb{C}^m$* , Proc. London Math. Soc. (3) **107** (2013), no. 2, 225–266.
- [74] G. Perelman, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, arXiv:math/0307245.
- [75] R. Schoen and S. T. Yau, *On the proof of the positive mass conjecture in general relativity*, Comm. Math. Phys. **65** (1979), no. 1, 45–76.
- [76] R. Schoen and S. T. Yau, *Positive scalar curvature and minimal hypersurface singularities*, arXiv:1704.05490.
- [77] A. Song, *Existence of infinitely many minimal hypersurfaces in closed manifolds*, arXiv:1806.08816.
- [78] M. B. Stenzel, *Ricci-flat metrics on the complexification of a compact rank one symmetric space*, Manuscripta Math. **80** (1993), no. 2, 151–163.